

HYDROXYPROPYL AMIDES FOR  
THE TREATMENT OF ALZHEIMER'S DISEASE

CROSS REFERENCE TO RELATED APPLICATIONS

5        This application claims priority from U.S. provisional application serial no. 60/515,908, filed October 30, 2003, which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

10        The invention relates to hydroxypropyl amides and to such compounds that are useful in the treatment of Alzheimer's disease and related diseases. More specifically, it relates to such compounds that are capable of inhibiting beta-secretase, an enzyme that cleaves amyloid precursor protein to  
15        produce amyloid beta peptide (A beta), a major component of the amyloid plaques found in the brains of Alzheimer's sufferers.

Background of the Invention

20        Alzheimer's disease (AD) is a progressive degenerative disease of the brain primarily associated with aging. Clinical presentation of AD is characterized by loss of memory, cognition, reasoning, judgment, and orientation. As the disease progresses, motor, sensory, and linguistic abilities are also affected until there is global impairment  
25        of multiple cognitive functions. These cognitive losses occur gradually, but typically lead to severe impairment and eventual death in the range of four to twelve years.

30        Alzheimer's disease is characterized by two major pathologic observations in the brain: neurofibrillary tangles and beta amyloid (or neuritic) plaques, comprised predominantly of an aggregate of a peptide fragment know as A beta. Individuals with AD exhibit characteristic beta-amyloid deposits in the brain (beta amyloid plaques) and in cerebral blood vessels (beta amyloid angiopathy) as well as  
35        neurofibrillary tangles. Neurofibrillary tangles occur not

only in Alzheimer's disease but also in other dementia-inducing disorders. On autopsy, large numbers of these lesions are generally found in areas of the human brain important for memory and cognition.

5       Smaller numbers of these lesions in a more restricted anatomical distribution are found in the brains of most aged humans who do not have clinical AD. Amyloidogenic plaques and vascular amyloid angiopathy also characterize the brains of individuals with Trisomy 21 (Down's Syndrome), Hereditary  
10 Cerebral Hemorrhage with Amyloidosis of the Dutch-Type (HCHWA-D), and other neurodegenerative disorders. Beta-amyloid is a defining feature of AD, now believed to be a causative precursor or factor in the development of disease. Deposition of A beta in areas of the brain responsible for cognitive  
15 activities is a major factor in the development of AD. Beta-amyloid plaques are predominantly composed of amyloid beta peptide (A beta, also sometimes designated betaA4). A beta peptide is derived by proteolysis of the amyloid precursor protein (APP) and is comprised of 39-42 amino acids. Several  
20 proteases called secretases are involved in the processing of APP.

Cleavage of APP at the N-terminus of the A beta peptide by beta-secretase and at the C-terminus by one or more gamma-secretases constitutes the beta-amyloidogenic pathway, i.e.  
25 the pathway by which A beta is formed. Cleavage of APP by alpha-secretase produces alpha-sAPP, a secreted form of APP that does not result in beta-amyloid plaque formation. This alternate pathway precludes the formation of A beta peptide. A description of the proteolytic processing fragments of APP  
30 is found, for example, in U.S. Patent Nos. 5,441,870; 5,721,130; and 5,942,400.

An aspartyl protease has been identified as the enzyme responsible for processing of APP at the beta-secretase cleavage site. The beta-secretase enzyme has been disclosed  
35 using varied nomenclature, including BACE, Asp, and Memapsin.

See, for example, Sinha et al., 1999, *Nature* 402:537-554 (p501) and published PCT application WO00/17369.

Several lines of evidence indicate that progressive cerebral deposition of beta-amyloid peptide (A beta) plays a seminal role in the pathogenesis of AD and can precede cognitive symptoms by years or decades. See, for example, Selkoe, 1991, *Neuron* 6:487. Release of A beta from neuronal cells grown in culture and the presence of A beta in cerebrospinal fluid (CSF) of both normal individuals and AD patients has been demonstrated. See, for example, Seubert et al., 1992, *Nature* 359:325-327.

It has been proposed that A beta peptide accumulates as a result of APP processing by beta-secretase, thus inhibition of this enzyme's activity is desirable for the treatment of AD. In vivo processing of APP at the beta-secretase cleavage site is thought to be a rate-limiting step in A beta production, and is thus a therapeutic target for the treatment of AD. See for example, Sabbagh, M., et al., 1997, *Alz. Dis. Rev.* 3, 1-19.

BACE1 knockout mice fail to produce A beta, and present a normal phenotype. When crossed with transgenic mice that over express APP, the progeny show reduced amounts of A beta in brain extracts as compared with control animals (Luo et al., 2001 *Nature Neuroscience* 4:231-232). This evidence further supports the proposal that inhibition of beta-secretase activity and reduction of A beta in the brain provides a therapeutic method for the treatment of AD and other beta amyloid disorders.


At present there are no effective treatments for halting, preventing, or reversing the progression of Alzheimer's disease. Therefore, there is an urgent need for pharmaceutical agents capable of slowing the progression of Alzheimer's disease and/or preventing it in the first place.

Compounds that are effective inhibitors of beta-secretase, that inhibit beta-secretase-mediated cleavage of

APP, that are effective inhibitors of A beta production, and/or are effective to reduce amyloid beta deposits or plaques, are needed for the treatment and prevention of disease characterized by amyloid beta deposits or plaques, such as AD.

# SUMMARY OF THE INVENTION

In one aspect, the invention provides compounds of formula (I):



and pharmaceutically acceptable salts thereof, wherein

$R_2$  is hydrogen, or

$R_2$  is  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_1-C_6 \text{ alkyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkenyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkynyl})-$  or  $(C_3-C_7 \text{ cycloalkyl})-$ , wherein each of said groups is optionally substituted with 1, 2, or 3  $R_2$  groups, wherein 1 or 2 methylene groups within said  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_1-C_6 \text{ alkyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkenyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkynyl})-$  or  $(C_3-C_7 \text{ cycloalkyl})-$  groups are optionally replaced with  $-(C=O)-$ ;

$R_2$  at each occurrence is independently halogen (in one aspect, F or Cl),  $-OH$ ,  $-SH$ ,  $-CN$ ,  $-CF_3$ ,  $-OCF_3$ ,  $C_1-C_6$  alkoxy,  $C_3-C_7$  cycloalkyl,  $C_3-C_7$  cycloalkoxy or  $-NR_{100}R_{101}$ ;

$R_{100}$  and  $R_{101}$  at each occurrence are independently H,  $C_1-C_6$  alkyl, phenyl,  $CO(C_1-C_6 \text{ alkyl})$  or  $SO_2C_1-C_6$  alkyl;

$X'$  is  $-(C=O)-$  or  $-(SO_2)-$ ;


$Y$  is absent or is  $-(CH_2)_n-$ , where  $n = 1, 2, \text{ or } 3$  and where up to 3 hydrogens of  $-(CH_2)_n-$  are optionally replaced with one, two or three substituents selected from

C<sub>1</sub>-C<sub>3</sub> alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF<sub>3</sub>,  
 C<sub>1</sub>-C<sub>3</sub> alkoxy, -COOH, -COO(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(COR)R', -  
 CONRR' or -NRR' where R and R' independently are -H  
 or C<sub>1</sub>-C<sub>10</sub> alkyl;

- 5 R<sub>1</sub> is H, -(CH<sub>2</sub>)<sub>1-2</sub>-S(O)<sub>0-2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), or  
 C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups  
 independently selected from halogen, -OH, =O, -SH,  
 -C≡N, -CF<sub>3</sub>, -COOR, C<sub>1</sub>-C<sub>3</sub> alkyl, -C<sub>1</sub>-C<sub>3</sub> alkoxy, amino,  
 monoalkylamino, dialkylamino, -CONRR', -N(R)C(O)R'-,  
 10 -OC(=O)-amino, -OC(=O)-monoalkylamino, and -OC(=O)-  
 dialkylamino or  
 C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally  
 substituted with 1, 2, or 3 groups independently  
 selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub>  
 15 alkoxy, amino, and mono- or dialkylamino, or  
 -C<sub>1</sub>-C<sub>6</sub> alkyl-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl where cycloalkyl can be  
 optionally substituted with C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -  
 OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, -O-phenyl, -CO<sub>2</sub>H, -  
 CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), or -NRR', or  
 20 aryl, heteroaryl, heterocyclyl, -C<sub>1</sub>-C<sub>6</sub> alkyl-aryl, -C<sub>1</sub>-C<sub>6</sub>  
 alkyl-heteroaryl, or -C<sub>1</sub>-C<sub>6</sub> alkyl-heterocyclyl, where  
 the ring portions of each are optionally substituted  
 with 1, 2, 3, or 4 of  
 halogen, -OH, -SH, -C≡N, -NRR', -CO<sub>2</sub>R, -  
 25 N(R)COR', or -N(R)SO<sub>2</sub>R', -C(=O)-(C<sub>1</sub>-C<sub>4</sub>) alkyl, -SO<sub>2</sub>-  
 amino, -SO<sub>2</sub>-mono or dialkylamino, -C(=O)-amino,  
 -C(=O)-mono or dialkylamino, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>) alkyl, or  
 -C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3  
 independently selected halogens, or  
 30 C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or  
 3 groups independently selected from halogen,  
 -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, -C<sub>1</sub>-C<sub>6</sub>  
 alkyl and mono- or dialkylamino, or

- $C_1$ - $C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, - $C_1$ - $C_3$  alkoxy, amino, mono- or dialkylamino and - $C_1$ - $C_3$  alkyl, or
- 5  $C_2$ - $C_{10}$  alkenyl or  $C_2$ - $C_{10}$  alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>,  $C_1$ - $C_3$  alkoxy, amino,  $C_1$ - $C_6$  alkyl and mono- or dialkylamino;
- 10 R and R' are independently -H or  $C_1$ - $C_{10}$  alkyl;  
 $R_3$  and  $R_3'$  at each occurrence are independently H,  $C_1$ - $C_6$  alkyl, -CO<sub>2</sub>- $C_1$ - $C_6$  alkyl, or -CO-O-(CH<sub>2</sub>)<sub>n</sub>-phenyl where n is 0, 1 or 2 and phenyl is optionally substituted with  $C_1$ - $C_6$  alkyl;
- 15  $R_4$  and  $R_5$  are independently H or  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>,  $C_1$ - $C_3$  alkoxy, and -NRR';

X is absent or is:



20

- $R_7$  is H,  $C_1$ - $C_6$  alkyl, -CO-O- $C_1$ - $C_6$  alkyl, or -CO-O-(CH<sub>2</sub>)<sub>n</sub>-phenyl where n is 0, 1 or 2 and phenyl is optionally substituted with  $C_1$ - $C_6$  alkyl, and wherein each  $C_1$ - $C_6$  alkyl is optionally independently substituted with
- 25 one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>,  $C_1$ - $C_6$  alkoxy, -NR<sub>102</sub>R'<sub>102</sub>,

Z is H,  $C_1$ - $C_6$  alkyl, CN, -O- $C_1$ - $C_6$  alkyl, or NO<sub>2</sub>;

R<sub>102</sub> and R'<sub>102</sub> independently are hydrogen, or

C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups that are independently halogen, or aryl, wherein aryl is optionally with 1 or 2 R<sub>125</sub> groups;

R<sub>125</sub> at each occurrence is independently halogen, amino, mono- or dialkylamino, -OH, -C≡N, -SO<sub>2</sub>-NH<sub>2</sub>, -SO<sub>2</sub>-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, -SO<sub>2</sub>-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, or -CO-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, or

C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, and mono- and dialkylamino, or

C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with one, two or three of halogen;

R<sub>6</sub> is -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-aryl, -[C(R<sub>255</sub>)(R<sub>260</sub>)]<sub>1-3</sub>-CO-N-(R<sub>255</sub>)<sub>2</sub>, -CH(aryl)<sub>2</sub>, -CH(heteroaryl)<sub>2</sub>, -CH(heterocyclyl)<sub>2</sub>, -CH(aryl)(heteroaryl), -(CH<sub>2</sub>)<sub>0-1</sub>-CH((CH<sub>2</sub>)<sub>0-6</sub>-OH)-(CH<sub>2</sub>)<sub>0-1</sub>-aryl, -(CH<sub>2</sub>)<sub>0-1</sub>-CH((CH<sub>2</sub>)<sub>0-6</sub>-OH)-(CH<sub>2</sub>)<sub>0-1</sub>-heteroaryl, -CH(-aryl or -heteroaryl)-CO-O(C<sub>1</sub>-C<sub>4</sub> alkyl), -CH(-CH<sub>2</sub>-OH)-CH(OH)-phenyl-NO<sub>2</sub>, -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-OH; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkenyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-aryl; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-cycloalkyl; -CH<sub>2</sub>-NH-CH<sub>2</sub>-CH(-O-CH<sub>2</sub>-CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-6</sub>-C(=NR<sub>235</sub>)(NR<sub>235</sub>R<sub>240</sub>), -(C<sub>2</sub>-C<sub>6</sub> alkenyl)-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl)-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-

heteroaryl,  $-(\text{CR}_{245}\text{R}_{250})_{1-4}-\text{C}(=\text{O})-\text{aryl}$ ,  $-(\text{CR}_{245}\text{R}_{250})_{1-4}-\text{C}(=\text{O})-\text{heteroaryl}$ , or

$\text{C}_1-\text{C}_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $\text{R}_{205}$ , cyclopentenyl,  $-\text{OC}=\text{ONR}_{235}\text{R}_{240}$ ,  $-\text{S}(=\text{O})_{0-2}(\text{C}_1-\text{C}_6 \text{ alkyl})$ ,  $-\text{SH}$ ,  $-\text{NR}_{235}\text{C}=\text{ONR}_{235}\text{R}_{240}$ ,  $-\text{C}=\text{ONR}_{235}\text{R}_{240}$ ,  $-\text{NR}_{235}-\text{C}(=\text{O})-\text{O}-\text{R}_{205}$ , and  $-\text{S}(=\text{O})_2\text{NR}_{235}\text{R}_{240}$ ,  $-\text{NR}_{235}\text{C}(=\text{O})-(\text{C}_1-\text{C}_6 \text{ alkyl})$ ,  $=\text{O}$ , or

$-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$  cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $\text{R}_{205}$ ,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$ ,  $-\text{CO}-\text{NH}_2$ ,  $-\text{CO}-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$  and  $-\text{CO}-\text{N}-(\text{C}_1-\text{C}_6 \text{ alkyl})(\text{C}_1-\text{C}_6 \text{ alkyl})$ , or

cyclopentyl, cyclohexyl, or cycloheptyl ring fused to aryl, heteroaryl, or heterocyclyl wherein one, two or three carbons of the cyclopentyl, cyclohexyl, or cycloheptyl is optionally replaced with a heteroatom independently selected from  $\text{NH}$ ,  $\text{NR}_{215}$ ,  $\text{O}$ , and  $\text{S}(=\text{O})_{0-2}$ , and wherein the cyclopentyl, cyclohexyl, or cycloheptyl group is optionally substituted with one or two groups that are independently  $\text{R}_{205}$ ,  $=\text{O}$ ,  $-\text{CO}-\text{NR}_{235}\text{R}_{240}$ , or  $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$ , or

$\text{C}_2-\text{C}_{10}$  alkenyl or  $\text{C}_2-\text{C}_{10}$  alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected  $\text{R}_{205}$  groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3  $\text{R}_{200}$ , and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 independently selected  $\text{R}_{210}$ , and each cycloalkyl is optionally substituted with 1 or 2  $\text{R}_{205}$  groups;

$\text{R}_{200}$  at each occurrence is independently selected from  $-\text{OH}$ ,  $-\text{NO}_2$ , halogen,  $-\text{CF}_3$ ,  $-\text{CO}_2\text{H}$ ,  $\text{C}\equiv\text{N}$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{220}\text{R}_{225}$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl})$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl})$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-\text{aryl}$ ,  $-(\text{CH}_2)_{0-4}-\text{CO}-\text{heteroaryl}$ , -



$(\text{CH}_2)_{0-4}\text{-CO-heterocyclyl}$ ,  $-(\text{CH}_2)_{0-4}\text{-CO-O-R}_{215}$ ,  $-(\text{CH}_2)_{0-4}\text{-SO}_2\text{-NR}_{220}\text{R}_{225}$ ,  $-(\text{CH}_2)_{0-4}\text{-SO-(C}_1\text{-C}_8\text{ alkyl)}$ ,  $-(\text{CH}_2)_{0-4}\text{-SO}_2\text{-(C}_1\text{-C}_{12}\text{ alkyl)}$ ,  $-(\text{CH}_2)_{0-4}\text{-SO}_2\text{-(C}_3\text{-C}_7\text{ cycloalkyl)}$ ,  $-(\text{CH}_2)_{0-4}\text{-N(H or R}_{215})\text{-CO-O-R}_{215}$ ,  $-(\text{CH}_2)_{0-4}\text{-N(H or R}_{215})\text{-CO-N(R}_{215})_2$ ,  $-(\text{CH}_2)_{0-4}\text{-N-CS-N(R}_{215})_2$ ,  $-(\text{CH}_2)_{0-4}\text{-N(-H or R}_{215})\text{-CO-R}_{220}$ ,  $-(\text{CH}_2)_{0-4}\text{-NR}_{220}\text{R}_{225}$ ,  $-(\text{CH}_2)_{0-4}\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$ ,  $-(\text{CH}_2)_{0-4}\text{-O-P(O)-(OR}_{240})_2$ ,  $-(\text{CH}_2)_{0-4}\text{-O-CO-N(R}_{215})_2$ ,  $-(\text{CH}_2)_{0-4}\text{-O-CS-N(R}_{215})_2$ ,  $-(\text{CH}_2)_{0-4}\text{-O-(R}_{215})$ ,  $-(\text{CH}_2)_{0-4}\text{-O-(R}_{215})\text{-COOH}$ ,  $-(\text{CH}_2)_{0-4}\text{-S-(R}_{215})$ ,  $-(\text{CH}_2)_{0-4}\text{-O-(C}_1\text{-C}_6\text{ alkyl optionally substituted with 1, 2, 3, or 5 -F, C}_3\text{-C}_7\text{ cycloalkyl)}$ ,  $-(\text{CH}_2)_{0-4}\text{-N(H or R}_{215})\text{-SO}_2\text{-R}_{220}$ ,  $-(\text{CH}_2)_{0-4}\text{-C}_3\text{-C}_7\text{ cycloalkyl}$ ,  $\text{C}_1\text{-C}_{10}\text{ alkyl optionally substituted with 1, 2, or 3 independently selected R}_{205}\text{ groups}$ ,  $\text{C}_2\text{-C}_{10}\text{ alkenyl and C}_2\text{-C}_{10}\text{ alkynyl, each of which is optionally substituted with 1 or 2 independently selected R}_{205}\text{ groups, wherein the aryl and heteroaryl groups at each occurrence are optionally substituted with 1, 2, or 3 groups that are independently R}_{205}, \text{R}_{210}, \text{or C}_1\text{-C}_6\text{ alkyl substituted with 1, 2, or 3 groups that are independently R}_{205}\text{ or R}_{210}, \text{and wherein the heterocyclyl group at each occurrence is optionally substituted with 1, 2, or 3 groups that are independently R}_{210}$ ;

$\text{R}_{205}$  at each occurrence is independently selected from  $\text{C}_1\text{-C}_6\text{ alkyl}$ , halogen,  $-\text{OH}$ ,  $-\text{COOH}$ ,  $-\text{O-phenyl}$ ,  $-\text{SH}$ ,  $-\text{S-C}_1\text{-C}_6\text{ alkyl}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $\text{C}_1\text{-C}_6\text{ alkoxy}$ ,  $\text{NH}_2$ ,  $\text{NH(C}_1\text{-C}_6\text{ alkyl)}$  or  $\text{N-(C}_1\text{-C}_6\text{ alkyl)(C}_1\text{-C}_6\text{ alkyl)}$ ;

$\text{R}_{210}$  at each occurrence is independently selected from halogen,  $\text{C}_1\text{-C}_6\text{ alkoxy}$ ,  $\text{C}_1\text{-C}_6\text{ haloalkoxy}$ ,  $-\text{NR}_{220}\text{R}_{225}$ ,  $\text{OH}$ ,  $\text{C}\equiv\text{N}$ ,  $-\text{CO-(C}_1\text{-C}_4\text{ alkyl)}$ ,  $-\text{SO}_2\text{-NR}_{235}\text{R}_{240}$ ,  $-\text{CO-NR}_{235}\text{R}_{240}$ ,  $-\text{C(=O)-(C}_1\text{-C}_6\text{ alkyl)-NR}_{235}\text{-C(=O)-O-R}_{205}$ ,  $-\text{C(=O)-(C}_1\text{-C}_4\text{ alkyl)-OH}$ ,  $-\text{C(=O)-(C}_1\text{-C}_6\text{ alkyl)-NR}_{235}\text{R}_{240}$ ,  $-\text{C(=O)-(C}_1\text{-C}_6\text{ alkyl)-imidazolyl}$ ,  $-\text{SO}_2\text{-(C}_1\text{-C}_4\text{ alkyl)}$ ,  $-\text{CO}_2\text{-(C}_1\text{-C}_4\text{ alkyl)}$ ,  $=\text{O}$ , or

C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or C<sub>3</sub>-C<sub>7</sub> cycloalkyl, each of which is optionally substituted with 1, 2, or 3 R<sub>205</sub> groups;

R<sub>215</sub> at each occurrence is independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, -(CH<sub>2</sub>)<sub>0-2</sub>-(aryl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, and -(CH<sub>2</sub>)<sub>0-2</sub>-(heteroaryl), -(CH<sub>2</sub>)<sub>0-2</sub>-(heterocyclyl), wherein

the aryl group at each occurrence is optionally substituted with 1, 2, or 3 groups that are independently R<sub>205</sub> or R<sub>210</sub>, and wherein

the heterocyclyl and heteroaryl groups at each occurrence are optionally substituted with 1, 2, or 3 independently selected R<sub>210</sub>;

R<sub>220</sub> and R<sub>225</sub> at each occurrence are independently selected from -H, -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and one triple bond, -aryl, -heteroaryl, and -heterocyclyl, and -C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with -OH, -NH<sub>2</sub> or halogen, wherein

the aryl, heterocyclyl and heteroaryl groups at each occurrence are optionally substituted with 1, 2, or 3 independently selected R<sub>270</sub> groups

R<sub>235</sub> and R<sub>240</sub> at each occurrence are independently H, or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sub>245</sub> and R<sub>250</sub> at each occurrence are independently selected from -H, halogen, -CF<sub>3</sub>, -OH, -NH<sub>2</sub>, -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, C<sub>1</sub>-C<sub>4</sub> alkylheteroaryl, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, -(CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, and phenyl; or

R<sub>245</sub> and R<sub>250</sub> are taken together with the carbon to which they are attached to form a carbocycle of 3, 4, 5, 6, or 7 carbon atoms, where one carbon atom is optionally replaced by a heteroatom selected from -O-, -S-, -SO<sub>2</sub>-, and -NR<sub>220</sub>-;

$R_{255}$  and  $R_{260}$  at each occurrence are independently selected from  
 -H,  $-(CH_2)_{1-2}-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$ ,  $-(C_1-C_4 \text{ alkyl})$ -aryl,  
 $-(C_1-C_4 \text{ alkyl})$ -heteroaryl,  $-(C_1-C_4 \text{ alkyl})$ -heterocyclyl, -  
 aryl, -heteroaryl, -heterocyclyl,  $-(CH_2)_{1-4}-R_{265}-(CH_2)_{0-4}$ -  
 5 aryl,  $-(CH_2)_{1-4}-R_{265}-(CH_2)_{0-4}$ -heteroaryl,  $-(CH_2)_{1-4}-R_{265}-(CH_2)_{0-4}$ -heterocyclyl, and

$C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl and  $-(CH_2)_{0-4}-C_3-C_7$   
 cycloalkyl, each of which is optionally substituted  
 with 1, 2, or 3 groups independently selected from  
 10  $R_{205}$ ,  $-COOH$ ,  $-COO(C_1-C_4 \text{ alkyl})$ ,  $-CO-NH_2$ ,  $-CO-NH(C_1-C_6 \text{ alkyl})$ ,  $-CO-N-(C_1-C_6 \text{ alkyl})(C_1-C_6 \text{ alkyl})$ , wherein  
 each aryl or phenyl is optionally substituted with 1, 2,  
 or 3 groups that are independently  $R_{205}$ ,  $R_{210}$ , or  
 $C_1-C_6$  alkyl substituted with 1, 2, or 3 groups that  
 15 are independently  $R_{205}$  or  $R_{210}$ , and wherein  
 each heterocyclyl is optionally substituted with 1, 2, 3,  
 or 4  $R_{210}$ ;

$R_{265}$  at each occurrence is independently -O-, -S- or  $-N(C_1-C_6 \text{ alkyl})$ -; and

20  $R_{270}$  at each occurrence is independently  $R_{205}$ , halogen  $C_1-C_6$   
 alkoxy,  $C_1-C_6$  haloalkoxy,  $NR_{235}R_{240}$ , -OH,  $-C\equiv N$ ,  $-CO-(C_1-C_4 \text{ alkyl})$ ,  
 $-SO_2-NR_{235}R_{240}$ ,  $-CO-NR_{235}R_{240}$ ,  $-SO_2-(C_1-C_4 \text{ alkyl})$ , =O,  
 or  
 $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl or  $-(CH_2)_{0-4}-C_3-C_7$   
 25 cycloalkyl, each of which is optionally substituted with  
 1, 2, or 3  $R_{205}$  groups.

The invention encompasses the compounds of formula (I)  
 shown above, pharmaceutical compositions containing the  
 30 compounds and methods employing such compounds or compositions  
 in the treatment of Alzheimer's disease and more specifically  
 compounds that are capable of inhibiting beta-secretase, an  
 enzyme that cleaves amyloid precursor protein to produce A-  
 beta peptide, a major component of the amyloid plaques found  
 35 in the brains of Alzheimer's sufferers.

Thus, the invention also provides methods for the treatment or prevention of Alzheimer's disease, mild cognitive impairment Down's syndrome, Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, cerebral amyloid angiopathy, other degenerative dementias, dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease comprising administration of a therapeutically effective amount of a compound or salt of formula I, to a patient in need thereof.

Preferably, the patient is a human.

More preferably, the disease is Alzheimer's disease.

More preferably, the disease is dementia.

The invention also provides pharmaceutical compositions comprising a compound or salt of formula I and at least one pharmaceutically acceptable carrier, solvent, adjuvant or diluent.

The invention also provides the use of a compound or salt according to formula I for the manufacture of a medicament.

The invention also provides the use of a compound or salt of formula I for the treatment or prevention of Alzheimer's disease, mild cognitive impairment Down's syndrome, Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, cerebral amyloid angiopathy, other degenerative dementias, dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, or diffuse Lewy body type of Alzheimer's disease.

The invention also provides compounds, pharmaceutical compositions, kits, and methods for inhibiting beta-secretase-mediated cleavage of amyloid precursor protein (APP). More particularly, the compounds, compositions, and methods of the

invention are effective to inhibit the production of A-beta peptide and to treat or prevent any human or veterinary disease or condition associated with a pathological form of A-beta peptide.

5       The compounds, compositions, and methods of the invention are useful for treating humans who have Alzheimer's Disease (AD), for helping prevent or delay the onset of AD, for treating patients with mild cognitive impairment (MCI), and preventing or delaying the onset of AD in those patients who  
10 would otherwise be expected to progress from MCI to AD, for treating Down's syndrome, for treating Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type, for treating cerebral beta-amyloid angiopathy and preventing its potential consequences such as single and recurrent lobar hemorrhages,  
15 for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, for treating dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy  
20 body type AD, and for treating frontotemporal dementias with parkinsonism (FTDP).

The compounds of the invention possess beta-secretase inhibitory activity. The inhibitory activities of the compounds of the invention is readily demonstrated, for  
25 example, using one or more of the assays described herein or known in the art.

Unless the substituents for a particular formula are expressly defined for that formula, they are understood to carry the definitions set forth in connection with the  
30 preceding formula to which the particular formula makes reference.


The invention also provides methods of preparing the compounds of the invention and the intermediates used in those methods.

DETAILED DESCRIPTION OF THE INVENTION


In one broad aspect, the invention encompasses compounds of formula I wherein X is




In another broad aspect, the invention encompasses compounds of formula I wherein X is



In another broad aspect, the invention encompasses compounds of formula I wherein X is



5 In another broad aspect, the invention encompasses compounds of formula I wherein X is




In another broad aspect, the invention encompasses compounds of formula I wherein X is



In another broad aspect, the invention encompasses compounds of formula I wherein X is



In another broad aspect, the invention encompasses compounds of formula I wherein X is



10 Preferred compounds of formula I include those of formula I-1, i.e., compounds of formula I wherein  $\text{R}_1$  and Y together form

15  $\text{C}_1$ - $\text{C}_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, =O, -SH, -C≡N, -CF<sub>3</sub>, -COOR,  $\text{C}_1$ - $\text{C}_3$  alkyl, - $\text{C}_1$ - $\text{C}_3$  alkoxy, amino, monoalkylamino, dialkylamino, -CONRR', -N(R)C(O)R',

-OC(=O)-amino, -OC(=O)-monoalkylamino, and -OC(=O)-dialkylamino or

C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, and mono- or dialkylamino, or

-C<sub>1</sub>-C<sub>6</sub> alkyl-(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl where cycloalkyl can be optionally substituted with C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, -O-phenyl, -CO<sub>2</sub>H, -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), or -NRR', or

aryl, heteroaryl, heterocyclyl, -C<sub>1</sub>-C<sub>6</sub> alkyl-aryl, -C<sub>1</sub>-C<sub>6</sub> alkyl-heteroaryl, or -C<sub>1</sub>-C<sub>6</sub> alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 of

halogen, -OH, -SH, -C≡N, -NRR', -CO<sub>2</sub>R, -N(R)COR', or -N(R)SO<sub>2</sub>R', -C(=O)-(C<sub>1</sub>-C<sub>4</sub>) alkyl, -SO<sub>2</sub>-amino, -SO<sub>2</sub>-mono or dialkylamino, -C(=O)-amino, -C(=O)-mono or dialkylamino, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>) alkyl, or -C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3

independently selected halogens, or C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, -C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino, or

C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, mono- or dialkylamino and -C<sub>1</sub>-C<sub>3</sub> alkyl, or

C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino.

Preferred compounds of formula I-1 also include those wherein:

$R_1$  and Y together form aryl, heteroaryl, heterocyclyl,  $-C_1-C_6$  alkyl-aryl,  $-C_1-C_6$  alkyl-heteroaryl, or  $-C_1-C_6$  alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-NO_2$ ,  $-NRR'$ ,  $-CO_2R$ ,  $-N(R)COR'$ , or  $-N(R)SO_2R'$ ,  $-C(=O)-(C_1-C_4)$  alkyl,  $-SO_2$ -amino,  $-SO_2$ -mono or dialkylamino,  $-C(=O)$ -amino,  $-C(=O)$ -mono or dialkylamino,  $-SO_2-(C_1-C_4)$  alkyl, or  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 groups which are independently selected from halogen, or  $C_3-C_7$  cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino,  $-C_1-C_6$  alkyl and mono- or dialkylamino, or  $C_1-C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $-C_1-C_3$  alkoxy, amino, mono- or dialkylamino and  $-C_1-C_3$  alkyl, or  $C_2-C_{10}$  alkenyl or  $C_2-C_{10}$  alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino,  $C_1-C_6$  alkyl and mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with oxo.

More preferred compounds of formula I-1 include those wherein

$R_1$  and Y together form  $-C_1-C_6$  alkyl-aryl,  $-C_1-C_6$  alkyl-heteroaryl, or  $-C_1-C_6$  alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected



from halogen, -OH, -SH, -C≡N, -NO<sub>2</sub>, -NRR', -CO<sub>2</sub>R, -N(R)COR', or -N(R)SO<sub>2</sub>R', -C(=O)-(C<sub>1</sub>-C<sub>4</sub>) alkyl, -SO<sub>2</sub>-amino, -SO<sub>2</sub>-mono or dialkylamino, -C(=O)-amino, -C(=O)-mono or dialkylamino, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>) alkyl, or  
5 C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 groups which are independently selected from halogen, or  
C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  
10 -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, -C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino, or  
C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, amino,  
15 mono- or dialkylamino and -C<sub>1</sub>-C<sub>3</sub> alkyl, or  
C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, C<sub>1</sub>-C<sub>6</sub> alkyl and  
20 mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with oxo.

Still more preferred compounds of formula I-1 include those wherein

25 R<sub>1</sub> and Y together are -(CH<sub>2</sub>)-aryl, where the aryl is optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, -OH, -SH, -C≡N, -NO<sub>2</sub>, -NRR', -CO<sub>2</sub>R, -N(R)COR', or -N(R)SO<sub>2</sub>R', -C(=O)-(C<sub>1</sub>-C<sub>4</sub>) alkyl, -SO<sub>2</sub>-amino, -SO<sub>2</sub>-mono or dialkylamino, -C(=O)-amino,  
30 -C(=O)-mono or dialkylamino, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>) alkyl, or C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 groups which are independently selected from halogen, or

C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, -C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino, or

5 C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, mono- or dialkylamino and -C<sub>1</sub>-C<sub>3</sub> alkyl, or

10 C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with  
15 oxo.

Yet more preferred compounds of formula I-1 include those wherein

R<sub>1</sub> and Y together are -CH<sub>2</sub>-phenyl where the phenyl ring is optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, C<sub>1</sub>-C<sub>4</sub> alkoxy, hydroxy, -NO<sub>2</sub>, and

20 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 1, 2, or 3 substituents independently selected from halogen, OH, SH, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub> alkyl), N-(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), C≡N, CF<sub>3</sub>.

25 Still more preferred compounds of formula I-1 include those wherein

R<sub>1</sub> and Y together are -CH<sub>2</sub>-phenyl where the phenyl ring is optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1</sub>-C<sub>2</sub> alkyl, C<sub>1</sub>-C<sub>2</sub> alkoxy, hydroxy, -CF<sub>3</sub>, and -NO<sub>2</sub>.

30 Preferred compounds of formula I-1 include those wherein R<sub>1</sub> and Y together are benzyl or 3,5-difluorobenzyl.

35 Preferred compounds of formula I-1 also include those wherein Y is absent and R<sub>1</sub> is hydrogen.

Preferred compounds of formula I-1 also include those wherein  $R_1$  and Y together form  $-(CH_2)_n$ -aryl, wherein n is 1, 2 or 3 and wherein 1, 2, or 3 hydrogens of  $-(CH_2)_n$ - are replaced with one, two or three groups independently selected from F, Cl, Br, I, OH,  $C_1$ - $C_3$  alkoxy,  $-N(COR)R'$ , and  $-NRR'$ . More preferably, n is 1.

Preferred compounds of formula I-1 also include those wherein  $R_1$  and Y together are  $-CH(OH)$ -aryl,  $-CH(halogen)$ -aryl,  $-CH(OMe)$ -aryl,  $-CH(OEt)$ -aryl,  $-CH(NH_2)$ -aryl,  $-CH(NMe_2)$ -aryl,  $-CH(NEt_2)$ -aryl,  $-CH(N(COMe)(Me))$ -aryl or  $-CH(N(COEt)(Et))$ -aryl.

Preferred compounds of formula I-1 also include those wherein  $R_1$  and Y together are  $-CH(OH)$ -difluorophenyl,  $-CH(halogen)$ -difluorophenyl,  $-CH(OMe)$ -difluorophenyl,  $-CH(OEt)$ -difluorophenyl,  $-CH(NH_2)$ -difluorophenyl,  $-CH(NMe_2)$ -difluorophenyl,  $-CH(NEt_2)$ -difluorophenyl,  $-CH(N(COMe)(Me))$ -difluorophenyl or  $-CH(N(COEt)(Et))$ -difluorophenyl.

Preferred compounds of formula I-1 also include those wherein  $R_1$  and Y together are  $-CH(OH)$ -difluorophenyl, more preferably  $-CH(OH)$ -3,5-difluorophenyl.

Preferred compounds of the formula I and formula I-1 include compounds of the formula I-2, i.e., compounds of the formula I or formula I-1 wherein:

$R_2$  is  $(C_3$ - $C_7$  cycloalkyl) $_{0-1}$ ( $C_1$ - $C_6$  alkyl)-,  $(C_3$ - $C_7$  cycloalkyl) $_{0-1}$ ( $C_2$ - $C_6$  alkenyl)-,  $(C_3$ - $C_7$  cycloalkyl) $_{0-1}$ ( $C_2$ - $C_6$  alkynyl)- or  $(C_3$ - $C_7$  cycloalkyl)-, wherein each of said groups is optionally substituted with 1, 2, or 3  $R_z$  groups;

$R_z$  at each occurrence is independently halogen,  $-OH$ ,  $-CN$ ,  $C_1$ - $C_6$  alkoxy,  $C_3$ - $C_7$  cycloalkyl,  $C_3$ - $C_7$  cycloalkoxy, or  $-NR_{100}R_{101}$ ;

$R_{100}$  and  $R_{101}$  are independently H,  $C_1$ - $C_6$  alkyl, phenyl,  $CO(C_1$ - $C_6$  alkyl) or  $SO_2C_1$ - $C_6$  alkyl.

Preferred compounds of formula I-2 include those wherein  $R_2$  is as defined above for formula I and  $X'$  is  $-(C=O)-$ . In an alternative embodiment,  $X'$  is  $-(C=O)-$ , and Z is H.

Preferred compounds of formula I-2 also include compounds wherein X' is  $-(C=O)-$ , and

R<sub>2</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1 or 2 groups independently selected from halogen (in one aspect, F or Cl), -OH, -SH, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxy or -NR<sub>100</sub>R<sub>101</sub>.

Preferred compounds of formula I-2 also include those wherein X' is  $-(C=O)-$ , and R<sub>2</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, or -NH<sub>2</sub>.

Preferred compounds of formula I-2 also include those wherein X' is  $-(C=O)-$ , and R<sub>2</sub> is methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, or 3-methylpentyl, any of which is optionally substituted with halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, or -NH<sub>2</sub>.

Preferred compounds of formula I-2 also include those wherein X' is  $-(C=O)-$ , and R<sub>2</sub> is methyl, ethyl, propyl, or isopropyl, any of which is optionally substituted with halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, or -NH<sub>2</sub>.

Preferred compounds of formula I-2 also include those wherein X' is  $-(C=O)-$ , and R<sub>2</sub> is methyl, ethyl or n-propyl.

Preferred compounds of formula I-2 also include those wherein X' is  $-(C=O)-$ , and R<sub>2</sub> is methyl.

Preferred compounds of the formula I, formula I-1, and formula I-2, include compounds of the formula I-3, i.e., compounds of the formula I, I-1, or I-2, wherein:

R<sub>3</sub> and R<sub>3</sub>' are independently H or C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferred compounds of the formula I-3 also include those wherein R<sub>3</sub> and R<sub>3</sub>' are both H.

Preferred compounds of the formula I, formula I-1, formula I-2, and formula I-3 include compounds of the formula I-4, i.e., compounds of the formula I, I-1, I-2, or I-3 wherein:

$R_4$  and  $R_5$  are independently H or  $C_1$ - $C_6$  alkyl, wherein  $C_1$ - $C_6$  alkyl is optionally substituted with one, two or three substituents selected from the group consisting of  $C_1$ - $C_3$  alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$ ,  $C_1$ - $C_3$  alkoxy, -NRR'.


Preferred compounds of the formula I-4 also include those wherein  $R_4$  and  $R_5$  are both H.

Preferred compounds of the formula I-4 also include those wherein  $R_4$  is H and  $R_5$  is  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents selected from the group consisting of  $C_1$ - $C_3$  alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C $\equiv$ N, -CF $_3$ ,  $C_1$ - $C_3$  alkoxy, -NRR'.

Preferred compounds of the formula I-4 also include those wherein  $R_4$  is H and  $R_5$  is  $C_1$ - $C_3$  alkyl optionally substituted with -OH.

Preferred compounds of the formula I-4 also include those wherein  $R_4$  is H and  $R_5$  is -CH $_3$ , -CH $_2$ CH $_3$ , -CH $_2$ CH $_2$ CH $_3$ , -CH $_2$ OH, or -CH $_2$ CH $_2$ OH.


Preferred compounds of the formula I, formula I-1, formula I-2, formula I-3, and formula I-4 include compounds of the formula I-5, i.e., compounds of the formula I, I-1, I-2, I-3 or I-4 wherein




$R_7$  is H,  $C_1$ - $C_6$  alkyl, -CO-O- $C_1$ - $C_6$  alkyl, or -CO-O-(CH $_2$ ) $_n$ -phenyl where  $n$  is 0, 1 or 2 and phenyl is optionally substituted with  $C_1$ - $C_6$  alkyl, and wherein each  $C_1$ - $C_6$  alkyl is optionally independently substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, -OH, -SH, -C $\equiv$ N, -CF $_3$ ,  $C_1$ - $C_6$  alkoxy, -NR $_{102}$ R' $_{102}$ ; and

Z is H,  $C_1$ - $C_6$  alkyl, CN, -O- $C_1$ - $C_6$  alkyl, or NO $_2$ .


Preferred compounds of the formula I-5 include compounds wherein:



Preferred compounds of the formula I-5 include compounds wherein:




Preferred compounds of the formula I-5 include compounds wherein:



R<sub>7</sub> is H; and

Z is -CN.

Preferred compounds of the formula I-5 include compounds wherein X is .

Preferred compounds of the formula I-5 include compounds wherein X is absent.

Preferred compounds of the formula I, formula I-1, formula I-2, formula I-3, formula I-4, and formula I-5 include compounds of the formula I-6, i.e., compounds of the formula I, I-1, I-2, I-3 I-4 or I-5 wherein

R<sub>6</sub> is -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-heterocyclyl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl-aryl, -[C(R<sub>255</sub>)(R<sub>260</sub>)]<sub>1-3</sub>-CO-N-(R<sub>255</sub>)<sub>2</sub>, -CH(aryl)<sub>2</sub>, -CH(heteroaryl)<sub>2</sub>, -CH(heterocyclyl)<sub>2</sub>, -CH(aryl)(heteroaryl), -(CH<sub>2</sub>)<sub>0-1</sub>-CH((CH<sub>2</sub>)<sub>0-6</sub>-OH)-(CH<sub>2</sub>)<sub>0-</sub>

<sub>1</sub>-aryl,  $-(CH_2)_{0-1}-CH((CH_2)_{0-6}-OH)-(CH_2)_{0-1}$ -heteroaryl,  
 $-CH(-aryl \text{ or } -heteroaryl)-CO-O(C_1-C_4 \text{ alkyl})$ ,  $-CH(-CH_2-OH)-CH(OH)-phenyl-NO_2$ ,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-OH$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkenyl})$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_0-C_6 \text{ alkyl})-aryl$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_0-C_6 \text{ alkyl})-cycloalkyl$ ;  $-CH_2-NH-CH_2-CH(-O-CH_2-CH_3)_2$ ,  
 $-(CH_2)_{0-6}-C(=NR_{235})(NR_{235}R_{240})$ ,  $-(C_2-C_6 \text{ alkenyl})$ -heteroaryl,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-O-(C_1-C_3 \text{ alkyl})-aryl$ ,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-(C_0-C_3 \text{ alkyl})-aryl$ ,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-(C_0-C_3 \text{ alkyl})-heteroaryl$ ,  $-(CR_{245}R_{250})_{1-4}-C(=O)-aryl$ ,  $-(CR_{245}R_{250})_{1-4}-C(=O)-heteroaryl$ , or  
 $C_1-C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $R_{205}$ , cyclopentenyl,  $-OC=ONR_{235}R_{240}$ ,  $-S(=O)_{0-2}(C_1-C_6 \text{ alkyl})$ ,  $-SH$ ,  $-NR_{235}C=ONR_{235}R_{240}$ ,  $-C=ONR_{235}R_{240}$ ,  $-NR_{235}-C(=O)-O-R_{205}$ , and  $-S(=O)_2NR_{235}R_{240}$ ,  $-NR_{235}C(=O)-(C_1-C_6 \text{ alkyl})$ ,  $=O$ , or  
 $-(CH_2)_{0-3}-(C_3-C_8)$  cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $R_{205}$ ,  $-CO_2H$ ,  $-CO_2-(C_1-C_4 \text{ alkyl})$ ,  $-CO-NH_2$ ,  $-CO-NH(C_1-C_6 \text{ alkyl})$  and  $-CO-N-(C_1-C_6 \text{ alkyl})(C_1-C_6 \text{ alkyl})$ , or  
 $C_2-C_{10}$  alkenyl or  $C_2-C_{10}$  alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected  $R_{205}$  groups, wherein  
each aryl and heteroaryl is optionally substituted with 1, 2, or 3  $R_{200}$ , and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 independently selected  $R_{210}$ , and each cycloalkyl is optionally substituted with 1 or 2  $R_{205}$  groups.

Preferred compounds of the formula I-6 include compounds wherein:

$R_6$  is  $-(CR_{245}R_{250})_{0-4}$ -aryl,  $-(CR_{245}R_{250})_{0-4}$ -heteroaryl,  $-(CR_{245}R_{250})_{0-4}$ -heterocyclyl,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-OH$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkenyl})$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ ;  $-(C_1-C_6 \text{ alkyl})-O-(C_0-C_6 \text{ alkyl})$ -aryl;  $-(C_1-C_6 \text{ alkyl})-O-(C_0-C_6 \text{ alkyl})$ -cycloalkyl;  $-CH_2-NH-CH_2-CH(-O-CH_2-CH_3)_2$ ,  $-(CH_2)_{0-6}-C(=NR_{235})(NR_{235}R_{240})$ ,  $-(C_2-C_6 \text{ alkenyl})$ -heteroaryl,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-O-(C_1-C_3 \text{ alkyl})$ -aryl,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-(C_0-C_3 \text{ alkyl})$ -aryl,  $-(CR_{245}R_{250})_{1-4}-N(R_{235})-C(=O)-(C_0-C_3 \text{ alkyl})$ -heteroaryl,  $-(CR_{245}R_{250})_{1-4}-C(=O)$ -aryl,  $-(CR_{245}R_{250})_{1-4}-C(=O)$ -heteroaryl, or

$C_1-C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $R_{205}$ , cyclopentenyl,  $-OC(=O)NR_{235}R_{240}$ ,  $-S(=O)_{0-2}(C_1-C_6 \text{ alkyl})$ ,  $-SH$ ,  $-NR_{235}C(=O)NR_{235}R_{240}$ ,  $-C(=O)NR_{235}R_{240}$ ,  $-NR_{235}-C(=O)-O-R_{205}$ , and  $-S(=O)_2NR_{235}R_{240}$ ,  $-NR_{235}C(=O)-(C_1-C_6 \text{ alkyl})$ ,  $=O$ , or

$-(CH_2)_{0-3}-(C_3-C_8 \text{ cycloalkyl})$  wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $R_{205}$ ,  $-CO_2H$ ,  $-CO_2-(C_1-C_4 \text{ alkyl})$ ,  $-CO-NH_2$ ,  $-CO-NH(C_1-C_6 \text{ alkyl})$  and  $-CO-N-(C_1-C_6 \text{ alkyl})(C_1-C_6 \text{ alkyl})$ , or

$C_2-C_{10}$  alkenyl or  $C_2-C_{10}$  alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected  $R_{205}$  groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3  $R_{200}$ , and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 independently selected  $R_{210}$ , and each cycloalkyl is optionally substituted with 1 or 2  $R_{205}$  groups.

Preferred compounds of the formula I-6 also include compounds wherein:

$R_6$  is  $-(CR_{245}R_{250})_{0-4}$ -aryl,  $-(CR_{245}R_{250})_{0-4}$ -heteroaryl,  $-(CR_{245}R_{250})_{0-4}$ -heterocyclyl,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkenyl})$ ;  $-(C_1-$



$C_6$  alkyl)-O-( $C_1$ - $C_6$  alkyl)-O-( $C_1$ - $C_6$  alkyl); -( $C_1$ - $C_6$  alkyl)-O-( $C_0$ - $C_6$  alkyl)-aryl; -( $C_1$ - $C_6$  alkyl)-O-( $C_0$ - $C_6$  alkyl)-cycloalkyl; -( $C_2$ - $C_6$  alkenyl)-heteroaryl;  
 -( $CR_{245}R_{250}$ )<sub>1-4</sub>-N( $R_{235}$ )-C(=O)-O-( $C_1$ - $C_3$  alkyl)-aryl;  
 5 -( $CR_{245}R_{250}$ )<sub>1-4</sub>-N( $R_{235}$ )-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-aryl;  
 -( $CR_{245}R_{250}$ )<sub>1-4</sub>-N( $R_{235}$ )-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-heteroaryl;  
 -( $CR_{245}R_{250}$ )<sub>1-4</sub>-C(=O)-aryl; -( $CR_{245}R_{250}$ )<sub>1-4</sub>-C(=O)-heteroaryl, or

$C_1$ - $C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halogen, -OH, -COOH, -O-phenyl, -C≡N, -CF<sub>3</sub>,  $C_1$ - $C_6$  alkoxy, NH<sub>2</sub>, NH( $C_1$ - $C_6$  alkyl) or N-( $C_1$ - $C_6$  alkyl)( $C_1$ - $C_6$  alkyl), cyclopentenyl, -C=ONR<sub>235</sub>R<sub>240</sub>, -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, -NR<sub>235</sub>-C(=O)-(C<sub>1</sub>- $C_6$  alkyl), and =O, or

15 -(CH<sub>2</sub>)<sub>0-3</sub>-(C<sub>3</sub>-C<sub>7</sub>) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $C_1$ - $C_6$  alkyl, halogen, -OH, -O-phenyl, -C≡N, -CF<sub>3</sub>,  $C_1$ - $C_6$  alkoxy, NH<sub>2</sub>, NH( $C_1$ - $C_6$  alkyl) or N-( $C_1$ - $C_6$  alkyl)( $C_1$ - $C_6$  alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH( $C_1$ - $C_6$  alkyl) and -CO-N-( $C_1$ - $C_6$  alkyl)( $C_1$ - $C_6$  alkyl), or

$C_2$ - $C_{10}$  alkenyl or  $C_2$ - $C_{10}$  alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected R<sub>205</sub> groups, wherein

25 each aryl and heteroaryl is optionally substituted with 1, 2, or 3 R<sub>200</sub>, and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 groups independently selected from -C(=O)-(C<sub>1</sub>- $C_6$  alkyl)-NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, -C(=O)-(C<sub>1</sub>-C<sub>4</sub> alkyl)-OH, -C(=O)-(C<sub>1</sub>- $C_6$  alkyl)-NR<sub>235</sub>R<sub>240</sub>, -C(=O)-(C<sub>1</sub>- $C_6$  alkyl)-imidazolyl, -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), and =O, and each cycloalkyl is optionally substituted with 1 or 2  $C_1$ - $C_6$  alkyl, halogen, -OH, -COOH, -O-phenyl, -C≡N, -CF<sub>3</sub>,

C<sub>1</sub>-C<sub>6</sub> alkoxy, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub> alkyl) or N-(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl) groups; and

R<sub>245</sub> and R<sub>250</sub> at each occurrence are independently selected from -H, halogen, -CF<sub>3</sub>, -OH, -NH<sub>2</sub>, -C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy.

Preferred compounds of the formula I-6 also include compounds wherein:

R<sub>6</sub> is -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-heterocyclyl, -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl)-aryl; -(C<sub>2</sub>-C<sub>6</sub> alkenyl)-heteroaryl; or

C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halogen, -OH, -O-phenyl, -C<sub>1</sub>-C<sub>6</sub> alkoxy, and -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, or

-(CH<sub>2</sub>)<sub>1-3</sub>-(C<sub>3</sub>-C<sub>7</sub>) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, halogen, -OH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, NH<sub>2</sub>, or

C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected R<sub>205</sub> groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3 of OH, -NO<sub>2</sub>, halogen, -CF<sub>3</sub>, -CO<sub>2</sub>H, C≡N, or C<sub>1</sub>-C<sub>6</sub> alkoxy, and wherein each heterocyclyl is optionally substituted with 1, 2, or 3 groups independently selected from -C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl)-NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, -C(=O)-(C<sub>1</sub>-C<sub>4</sub> alkyl)-OH, and -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl); and

R<sub>245</sub> and R<sub>250</sub> at each occurrence are independently selected from -H, halogen, -CF<sub>3</sub>, -OH, -NH<sub>2</sub>, -C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy.

Preferred compounds of the formula I-6 also include compounds wherein:

$R_6$  is  $-(CH_2)_{1-4}$ -aryl,  $-(CH_2)_{1-4}$ -heteroaryl,  $-(CH_2)_{1-4}$ -heterocyclyl,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$ -aryl;  $-(C_2-C_6 \text{ alkenyl})$ -heteroaryl; or

$C_1-C_6$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halogen,  $-OH$ ,  $-O$ -phenyl,  $-C_1-C_6$  alkoxy, and  $-NR_{235}-C(=O)-O-R_{205}$ , or

$-(CH_2)_{1-3}$ -cyclopentyl wherein the cyclopentyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $C_1-C_6$  alkyl, halogen,  $-OH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_6$  alkoxy,  $NH_2$ , or

$C_2-C_6$  alkenyl optionally substituted with 1, 2, or 3 independently selected  $R_{205}$  groups, wherein each aryl and heteroaryl is optionally substituted with 1 or 2 of halogen,  $-CF_3$ ,  $C\equiv N$ , or  $C_1-C_6$  alkoxy.

Preferred compounds of the formula I, formula I-1, formula I-2, formula I-3, formula I-4, formula I-5 and formula I-6 include compounds of the formula I-7, i.e., compounds of the formula I, I-1, I-2, I-3, I-4, I-5 or I-6 wherein:

$R_1$  and  $Y$  together are  $-(CH_2)$ -aryl, where the aryl is optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-NO_2$ ,  $-NRR'$ ,  $-CO_2R$ ,  $-N(R)COR'$ , or  $-N(R)SO_2R'$ ,  $-C(=O)-(C_1-C_4 \text{ alkyl})$ ,  $-SO_2$ -amino,  $-SO_2$ -mono or dialkylamino,  $-C(=O)$ -amino,  $-C(=O)$ -mono or dialkylamino,  $-SO_2-(C_1-C_4 \text{ alkyl})$ ;

$X'$  is  $-(C=O)-$ ;

$R_2$  is  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_1-C_6 \text{ alkyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkenyl})-$ ,  $(C_3-C_7 \text{ cycloalkyl})_{0-1}(C_2-C_6 \text{ alkynyl})-$  or  $(C_3-C_7 \text{ cycloalkyl})-$ , wherein each of said groups is optionally substituted with 1, 2, or 3  $R_z$  groups;

$R_z$  at each occurrence is independently halogen,  $-OH$ ,  $-CN$ ,  $C_1-C_6$  alkoxy,  $C_3-C_7$  cycloalkyl,  $C_3-C_7$  cycloalkoxy, or  $-NR_{100}R_{101}$ ;

R<sub>100</sub> and R<sub>101</sub> are independently H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, CO(C<sub>1</sub>-C<sub>6</sub> alkyl) or SO<sub>2</sub>C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sub>3</sub> and R<sub>3</sub>' are both H;

R<sub>4</sub> and R<sub>5</sub> are both H;


5 R<sub>6</sub> is -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl, -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-OH; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkenyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-aryl; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-cycloalkyl; -CH<sub>2</sub>-NH-CH<sub>2</sub>-CH(-O-CH<sub>2</sub>-CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-6</sub>-C(=NR<sub>235</sub>)(NR<sub>235</sub>R<sub>240</sub>), -(C<sub>2</sub>-C<sub>6</sub> alkenyl)-heteroaryl, - (CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl)-aryl, - (CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-aryl, - (CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-heteroaryl, 15 - (CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-C(=O)-aryl, - (CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-C(=O)-heteroaryl, or

C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, cyclopentenyl, -OC=ONR<sub>235</sub>R<sub>240</sub>, -S(=O)<sub>0-2</sub>(C<sub>1</sub>-C<sub>6</sub> 20 alkyl), -SH, -NR<sub>235</sub>C=ONR<sub>235</sub>R<sub>240</sub>, -C=ONR<sub>235</sub>R<sub>240</sub>, -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, and -S(=O)<sub>2</sub>NR<sub>235</sub>R<sub>240</sub>, -NR<sub>235</sub>C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), =O, or

25 -(CH<sub>2</sub>)<sub>0-3</sub>-(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -CO<sub>2</sub>H, -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH(C<sub>1</sub>-C<sub>6</sub> alkyl) and -CO-N-(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), or

30 C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected R<sub>205</sub> groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3 R<sub>200</sub>, and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 independently selected R<sub>210</sub>, and each cycloalkyl is 35 optionally substituted with 1 or 2 R<sub>205</sub> groups;




R<sub>7</sub> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, -CO-O-C<sub>1</sub>-C<sub>6</sub> alkyl, or -CO-O-(CH<sub>2</sub>)<sub>n</sub>-phenyl where n is 0, 1 or 2 and phenyl is optionally substituted with C<sub>1</sub>-C<sub>6</sub> alkyl, and wherein each C<sub>1</sub>-C<sub>6</sub> alkyl is optionally independently substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, -NR<sub>102</sub>R'<sub>102</sub>; and

Z is H, C<sub>1</sub>-C<sub>6</sub> alkyl, CN, -O-C<sub>1</sub>-C<sub>6</sub> alkyl, or NO<sub>2</sub>.

Preferred compounds of the formula I-7 include compounds

of the formula I-7-a:




I-7-a

wherein

B<sub>1</sub> and B<sub>2</sub> are independently halogen, -OH, -SH, -C≡N, -NO<sub>2</sub>, -NRR', or -CO<sub>2</sub>R;

R<sub>2</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1 or 2 groups independently selected from halogen (in one aspect, F or Cl), -OH, -SH, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxy or -NR<sub>100</sub>R<sub>101</sub>;



R<sub>7</sub> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, -CO-O-C<sub>1</sub>-C<sub>6</sub> alkyl, or -CO-O-(CH<sub>2</sub>)<sub>n</sub>-phenyl where n is 0, 1 or 2 and phenyl is optionally substituted with C<sub>1</sub>-C<sub>6</sub> alkyl, and wherein each C<sub>1</sub>-C<sub>6</sub> alkyl is optionally independently substituted with

one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, -NR<sub>102</sub>R'<sub>102</sub>;

Z is H, C<sub>1</sub>-C<sub>6</sub> alkyl, CN, -O-C<sub>1</sub>-C<sub>6</sub> alkyl, or NO<sub>2</sub>;

5 R<sub>5</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, -NRR'; and

10 R<sub>6</sub> is -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocyclyl, -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-OH; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkenyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl); -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-aryl; -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl)-cycloalkyl; -CH<sub>2</sub>-NH-CH<sub>2</sub>-CH(-O-CH<sub>2</sub>-CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-6</sub>-C(=NR<sub>235</sub>)(NR<sub>235</sub>R<sub>240</sub>), -(C<sub>2</sub>-C<sub>6</sub> alkenyl)-heteroaryl, 15 -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl)-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-N(R<sub>235</sub>)-C(=O)-(C<sub>0</sub>-C<sub>3</sub> alkyl)-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-C(=O)-aryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>1-4</sub>-C(=O)-heteroaryl, or 20

C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, cyclopentenyl, -OC=ONR<sub>235</sub>R<sub>240</sub>, -S(=O)<sub>0-2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -SH, -NR<sub>235</sub>C=ONR<sub>235</sub>R<sub>240</sub>, -C=ONR<sub>235</sub>R<sub>240</sub>, -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, and -S(=O)<sub>2</sub>NR<sub>235</sub>R<sub>240</sub>, -NR<sub>235</sub>C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), =O, or 25

-(CH<sub>2</sub>)<sub>0-3</sub>-(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -CO<sub>2</sub>H, -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH(C<sub>1</sub>-C<sub>6</sub> alkyl) and -CO-N-(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), or 30


C<sub>2</sub>-C<sub>10</sub> alkenyl or C<sub>2</sub>-C<sub>10</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected R<sub>205</sub> groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3  $R_{200}$ , and wherein each heterocyclyl is optionally substituted with 1, 2, 3, or 4 independently selected  $R_{210}$ , and each cycloalkyl is optionally substituted with 1 or 2  $R_{205}$  groups.

Preferred compounds of the formula I-7-a include compounds wherein

$B_1$  and  $B_2$  are halogen;

$R_2$  is  $C_1$ - $C_6$  alkyl optionally substituted with halogen, -OH, -SH, - $C\equiv N$ , - $CF_3$ , - $C_1$ - $C_3$  alkoxy, or - $NH_2$ ;

X is ;

$R_5$  is  $C_1$ - $C_3$  alkyl optionally substituted with -OH; and

$R_6$  is -( $CR_{245}R_{250}$ )<sub>1-4</sub>-aryl, -( $CR_{245}R_{250}$ )<sub>1-4</sub>-heteroaryl, -( $CR_{245}R_{250}$ )<sub>1-4</sub>-heterocyclyl, -( $C_1$ - $C_6$  alkyl)-O-( $C_1$ - $C_3$  alkyl)-aryl; -( $C_2$ - $C_6$  alkenyl)-heteroaryl; or

$C_1$ - $C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halogen, -OH, -O-phenyl, - $C_1$ - $C_6$  alkoxy, and - $NR_{235}$ -C(=O)-O- $R_{205}$ , or


-( $CH_2$ )<sub>1-3</sub>-( $C_3$ - $C_7$ ) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of  $C_1$ - $C_6$  alkyl, halogen, -OH, - $C\equiv N$ , - $CF_3$ ,  $C_1$ - $C_6$  alkoxy,  $NH_2$ , or

$C_2$ - $C_{10}$  alkenyl or  $C_2$ - $C_{10}$  alkynyl, each of which is optionally substituted with 1, 2, or 3 independently selected  $R_{205}$  groups, wherein

each aryl and heteroaryl is optionally substituted with 1, 2, or 3 of OH, - $NO_2$ , halogen, - $CF_3$ , - $CO_2H$ ,  $C\equiv N$ , or  $C_1$ - $C_6$  alkoxy, and wherein each heterocyclyl is optionally substituted with 1, 2, or 3 groups independently selected from -C(=O)-( $C_1$ - $C_6$  alkyl)- $NR_{235}$ -C(=O)-O- $R_{205}$ , -C(=O)-( $C_1$ - $C_4$  alkyl)-OH, and - $CO_2$ -( $C_1$ - $C_4$  alkyl); and

R<sub>245</sub> and R<sub>250</sub> at each occurrence are independently selected from -H, halogen, -CF<sub>3</sub>, -OH, -NH<sub>2</sub>, -C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy.

Preferred compounds of formula I-7-a include compounds of  
5 formula I-7-b:



wherein,

B<sub>1</sub> and B<sub>2</sub> are halogen;

X is

R<sub>5</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with -OH; and

10 R<sub>6</sub> is -(CH<sub>2</sub>)<sub>1-4</sub>-aryl, -(CH<sub>2</sub>)<sub>1-4</sub>-heteroaryl, -(CH<sub>2</sub>)<sub>1-4</sub>-heterocyclyl, -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl)-aryl; -(C<sub>2</sub>-C<sub>6</sub> alkenyl)-heteroaryl; or

15 C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of halogen, -OH, -O-phenyl, -C<sub>1</sub>-C<sub>6</sub> alkoxy, and -NR<sub>235</sub>-C(=O)-O-R<sub>205</sub>, or

20 -(CH<sub>2</sub>)<sub>1-3</sub>-cyclopentyl wherein the cyclopentyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, halogen, -OH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, NH<sub>2</sub>, or


C<sub>2</sub>-C<sub>6</sub> alkenyl optionally substituted with 1, 2, or 3 independently selected R<sub>205</sub> groups, wherein

each aryl and heteroaryl is optionally substituted with 1, or 2 of halogen, -CF<sub>3</sub>, C≡N, or C<sub>1</sub>-C<sub>6</sub> alkoxy.

25



In another aspect, the invention provides intermediates of formula 23:



wherein  $R_1$  and  $R_2$  are as defined above for compounds of formula I.

Preferred intermediates of formula 23 include those wherein

$R_1$  is  $-C_1-C_6$  alkyl-aryl,  $-C_1-C_6$  alkyl-heteroaryl, or  $-C_1-C_6$  alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-NO_2$ ,  $-NRR'$ ,  $-CO_2R$ ,  $-N(R)COR'$ , or  $-N(R)SO_2R'$ ,  $-C(=O)-(C_1-C_4)$  alkyl,  $-SO_2$ -amino,  $-SO_2$ -mono or dialkylamino,  $-C(=O)$ -amino,  $-C(=O)$ -mono or dialkylamino,  $-SO_2-(C_1-C_4)$  alkyl, or

$C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 groups which are independently selected from halogen, or

$C_3-C_7$  cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino,  $-C_1-C_6$  alkyl and mono- or dialkylamino, or

$C_1-C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $-C_1-C_3$  alkoxy, amino, mono- or dialkylamino and  $-C_1-C_3$  alkyl, or

$C_2-C_{10}$  alkenyl or  $C_2-C_{10}$  alkynyl each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-OH$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino,  $C_1-C_6$  alkyl and

mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with oxo; and

R<sub>2</sub> is hydrogen, or

5 R<sub>2</sub> is (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl)-, (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>2</sub>-C<sub>6</sub> alkenyl)-, (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>2</sub>-C<sub>6</sub> alkynyl)- or (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)-, wherein each of said groups is optionally substituted with 1, 2, or 3 R<sub>z</sub> groups, wherein 1 or 2 methylene groups within said (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl)-, (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>2</sub>-C<sub>6</sub> alkenyl)-, (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)<sub>0-1</sub>(C<sub>2</sub>-C<sub>6</sub> alkynyl)- or (C<sub>3</sub>-C<sub>7</sub> cycloalkyl)- groups are optionally replaced with -(C=O)-;

R<sub>z</sub> at each occurrence is independently halogen (in one aspect, F or Cl), -OH, -SH, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxy or -NR<sub>100</sub>R<sub>101</sub>;

R<sub>100</sub> and R<sub>101</sub> at each occurrence are independently H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, CO(C<sub>1</sub>-C<sub>6</sub> alkyl) or SO<sub>2</sub>C<sub>1</sub>-C<sub>6</sub> alkyl;


20 Preferred intermediates of formula 23 also include those wherein

R<sub>1</sub> is -CH<sub>2</sub>-phenyl where the phenyl ring is optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, C<sub>1</sub>-C<sub>4</sub> alkoxy, hydroxy, -NO<sub>2</sub>, and

25 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 1, 2, or 3 substituents independently selected from halogen, OH, SH, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub> alkyl), N-(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), C≡N, CF<sub>3</sub>; and


30 R<sub>2</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1 or 2 groups independently selected from halogen (in one aspect, F or Cl), -OH, -SH, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxy or -NR<sub>100</sub>R<sub>101</sub>.

The invention also provides intermediates of formula 19:




wherein  $R_1$  is as defined above for compounds of formula I.

The invention further provides intermediates of formula 20:



wherein  $R_1$  is as defined above for compounds of formula I.

The invention also provides intermediates of formula 21:



wherein  $R_1$  is as defined above for compounds of formula I.

The invention also provides methods for treating a patient who has, or in preventing a patient from getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, or diffuse Lewy body type of Alzheimer's disease and who is in need of such treatment which includes administration of a therapeutically

effective amount of a compound of formula (I) or a pharmaceutically acceptable salts thereof.

In an embodiment, this method of treatment can be used where the disease is Alzheimer's disease.

5 In an embodiment, this method of treatment can help prevent or delay the onset of Alzheimer's disease.

In an embodiment, this method of treatment can be used where the disease is mild cognitive impairment.

10 In an embodiment, this method of treatment can be used where the disease is Down's syndrome.

In an embodiment, this method of treatment can be used where the disease is Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type.

15 In an embodiment, this method of treatment can be used where the disease is cerebral amyloid angiopathy.

In an embodiment, this method of treatment can be used where the disease is degenerative dementias.

20 In an embodiment, this method of treatment can be used where the disease is diffuse Lewy body type of Alzheimer's disease.

In an embodiment, this method of treatment can treat an existing disease.

In an embodiment, this method of treatment can prevent a disease from developing.

25 In an embodiment, this method of treatment can employ therapeutically effective amounts: for oral administration from about 0.1 mg/day to about 1,000 mg/day; for parenteral, sublingual, intranasal, intrathecal administration from about 0.5 to about 100 mg/day; for depo administration and implants  
30 from about 0.5 mg/day to about 50 mg/day; for topical administration from about 0.5 mg/day to about 200 mg/day; for rectal administration from about 0.5 mg to about 500 mg.

In an embodiment, this method of treatment can employ therapeutically effective amounts: for oral administration

from about 1 mg/day to about 100 mg/day; and for parenteral administration from about 5 to about 50 mg daily.

In an embodiment, this method of treatment can employ therapeutically effective amounts for oral administration from  
5 about 5 mg/day to about 50 mg/day.

The invention also includes pharmaceutical compositions which include a compound of formula (I) or a pharmaceutically acceptable salts thereof.

The invention also includes the use of a compound of  
10 formula (I) or pharmaceutically acceptable salts thereof for the manufacture of a medicament for use in treating a patient who has, or in preventing a patient from getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's  
15 disease, for treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-  
20 Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia  
25 associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease and who is in need of such treatment.

In an embodiment, this use of a compound of formula (I)  
30 can be employed where the disease is Alzheimer's disease.

In an embodiment, this use of a compound of formula (I) can help prevent or delay the onset of Alzheimer's disease.

In an embodiment, this use of a compound of formula (I) can be employed where the disease is mild cognitive  
35 impairment.

In an embodiment, this use of a compound of formula (I) can be employed where the disease is Down's syndrome.

In an embodiment, this use of a compound of formula (I) can be employed where the disease is Hereditary Cerebral  
5 Hemorrhage with Amyloidosis of the Dutch-Type.

In an embodiment, this use of a compound of formula (I) can be employed where the disease is cerebral amyloid angiopathy.

In an embodiment, this use of a compound of formula (I)  
10 can be employed where the disease is degenerative dementias.

In an embodiment, this use of a compound of formula (I) can be employed where the disease is diffuse Lewy body type of Alzheimer's disease.

In an embodiment, this use of a compound employs a  
15 pharmaceutically acceptable salt selected from the group consisting of salts of the following acids hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, citric, methanesulfonic,  $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$  where  $n$  is 0 thru 4,  $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$  where  $n$  is as defined above,  $\text{HOOC}-\text{CH}=\text{CH}-\text{COOH}$ , and  
20 phenyl-COOH.

The invention also includes methods for inhibiting beta-secretase activity, for inhibiting cleavage of amyloid precursor protein (APP), in a reaction mixture, at a site between Met596 and Asp597, numbered for the APP-695 amino acid  
25 isotype, or at a corresponding site of an isotype or mutant thereof; for inhibiting production of amyloid beta peptide (A beta) in a cell; for inhibiting the production of beta-amyloid plaque in an animal; and for treating or preventing a disease characterized by beta-amyloid deposits in the brain. These  
30 methods each include administration of a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salts thereof.

The invention also includes a method for inhibiting beta-secretase activity, including exposing said beta-secretase to

an effective inhibitory amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of  
5 less than 50 micromolar.

In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of 10 micromolar or less.

In an embodiment, this method employs a compound that  
10 inhibits 50% of the enzyme's activity at a concentration of 1 micromolar or less.

In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of 10 nanomolar or less.

15 In an embodiment, this method includes exposing said beta-secretase to said compound *in vitro*.

In an embodiment, this method includes exposing said beta-secretase to said compound in a cell.

In an embodiment, this method includes exposing said  
20 beta-secretase to said compound in a cell in an animal.

In an embodiment, this method includes exposing said beta-secretase to said compound in a human.

The invention also includes a method for inhibiting cleavage of amyloid precursor protein (APP), in a reaction  
25 mixture, at a site between Met596 and Asp597, numbered for the APP-695 amino acid isotype; or at a corresponding site of an isotype or mutant thereof, including exposing said reaction mixture to an effective inhibitory amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

30 In an embodiment, this method employs a cleavage site: between Met652 and Asp653, numbered for the APP-751 isotype; between Met 671 and Asp 672, numbered for the APP-770 isotype; between Leu596 and Asp597 of the APP-695 Swedish Mutation; between Leu652 and Asp653 of the APP-751 Swedish Mutation; or  
35 between Leu671 and Asp672 of the APP-770 Swedish Mutation.

In an embodiment, this method exposes said reaction mixture *in vitro*.

In an embodiment, this method exposes said reaction mixture in a cell.

5 In an embodiment, this method exposes said reaction mixture in an animal cell.

In an embodiment, this method exposes said reaction mixture in a human cell.

10 The invention also includes a method for inhibiting production of amyloid beta peptide (A beta) in a cell, including administering to said cell an effective inhibitory amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

15 In an embodiment, this method includes administering to an animal.

In an embodiment, this method includes administering to a human.

20 The invention also includes a method for inhibiting the production of beta-amyloid plaque in an animal, including administering to said animal an effective inhibitory amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

In an embodiment, this method includes administering to a human.

25 The invention also includes a method for treating or preventing a disease characterized by beta-amyloid deposits in the brain including administering to a patient an effective therapeutic amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

30 In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of less than 50 micromolar.

In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of 10  
35 micromolar or less.



In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of 1 micromolar or less.

5 In an embodiment, this method employs a compound that inhibits 50% of the enzyme's activity at a concentration of 10 nanomolar or less.

In an embodiment, this method employs a compound at a therapeutic amount in the range of from about 0.1 to about 1000 mg/day.

10 In an embodiment, this method employs a compound at a therapeutic amount in the range of from about 15 to about 1500 mg/day.

In an embodiment, this method employs a compound at a therapeutic amount in the range of from about 1 to about 100  
15 mg/day.

In an embodiment, this method employs a compound at a therapeutic amount in the range of from about 5 to about 50 mg/day.

In an embodiment, this method can be used where said  
20 disease is Alzheimer's disease.

In an embodiment, this method can be used where said disease is Mild Cognitive Impairment, Down's Syndrome, or Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type.

25 The invention also includes a composition including beta-secretase complexed with a compound of formula (I), or a pharmaceutically acceptable salt thereof.

The invention also includes a method for producing a beta-secretase complex including exposing beta-secretase to a  
30 compound of formula (I), or a pharmaceutically acceptable salt thereof, in a reaction mixture under conditions suitable for the production of said complex.

In an embodiment, this method employs exposing *in vitro*.

In an embodiment, this method employs a reaction mixture  
35 that is a cell.

The invention also includes a component kit including component parts capable of being assembled, in which at least one component part includes a compound of formula I enclosed in a container.

5        In an embodiment, this component kit includes lyophilized compound, and at least one further component part includes a diluent.

10        The invention also includes a container kit including a plurality of containers, each container including one or more unit dose of a compound of formula (I):, or a pharmaceutically acceptable salt thereof.

      In an embodiment, this container kit includes each container adapted for oral delivery and includes a tablet, gel, or capsule.

15        In an embodiment, this container kit includes each container adapted for parenteral delivery and includes a depot product, syringe, ampoule, or vial.

20        In an embodiment, this container kit includes each container adapted for topical delivery and includes a patch, medipad, ointment, or cream.

25        The invention also includes an agent kit including a compound of formula (I), or a pharmaceutically acceptable salt thereof; and one or more therapeutic agent selected from the group consisting of an antioxidant, an anti-inflammatory, a gamma secretase inhibitor, a neurotrophic agent, an acetyl cholinesterase inhibitor, a statin, an A beta peptide, and an anti-A beta antibody.

30        The invention also includes a composition including a compound of formula (I), or a pharmaceutically acceptable salt thereof; and an inert diluent or edible carrier.

      In an embodiment, this composition includes a carrier that is an oil.

      The invention also includes a composition including: a compound of formula (I), or a pharmaceutically acceptable salt


thereof; and a binder, excipient, disintegrating agent, lubricant, or gildant.

The invention also includes a composition including a compound of formula (I), or a pharmaceutically acceptable salt thereof; disposed in a cream, ointment, or patch.


The invention provides compounds of formula (I) that are useful in treating and preventing Alzheimer's disease. The compounds of the invention can be prepared by one skilled in the art based only on knowledge of the compound's chemical structure. The chemistry for the preparation of the compounds of this invention is known to those skilled in the art. In fact, there is more than one process to prepare the compounds of the invention. Specific examples of methods of preparation can be found in the art. For examples, see *J. Org. Chem.* **1998**, 63, 4898-4906; *J. Org. Chem.* **1997**, 62, 9348-9353; *J. Org. Chem.* **1996**, 61, 5528-5531; *J. Med. Chem.* **1993**, 36, 320-330; *J. Am. Chem. Soc.* **1999**, 121, 1145-1155; *J. Org. Chem.*, 1992, vol 57, 5692-5700; *J. Org. Chem.* 1994, vol 59, 4040-4041, and especially *J. Org. Chem.* 1993, vol 58, 1025-1029; and references cited therein. See also U.S. Patent Nos. 6,150,530, 5,892,052, 5,696,270, and 5,362,912, which are incorporated herein by reference, and references cited therein.

Examples of various processes that can be used to prepare the compounds of the invention are set forth in Schemes 1-7.


Scheme 1.




Scheme 2




Scheme 3



## SCHEME 4




Scheme 5






Scheme 6.



Scheme 7




The protection of amines is conducted, where appropriate,  
 5 by methods known to those skilled in the art. Amino protecting  
 groups are known to those skilled in the art. See for  
 example, "Protecting Groups in Organic Synthesis", John Wiley

and sons, New York, N.Y., 1981, Chapter 7; "Protecting Groups in Organic Chemistry", Plenum Press, New York, N.Y., 1973, Chapter 2. When the amino protecting group is no longer needed, it is removed by methods known to those skilled in the art. By definition the amino protecting group must be readily removable. A variety of suitable methodologies are known to those skilled in the art; see also T.W. Green and P.G.M. Wuts in "Protective Groups in Organic Chemistry, John Wiley and Sons, 1991. Suitable amino protecting groups include t-butoxycarbonyl, benzyl-oxycarbonyl, formyl, trityl, phthalimido, trichloro-acetyl, chloroacetyl, bromoacetyl, iodoacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluy)prop-2-yloxy-carbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycabonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluy)sulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)-ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxy-carbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxyl)benzyloxycarbonyl, isobronnyloxycarbonyl, 1-piperidyloxycarbonyl, 9-fluoroenylmethyl carbonate, -CH-CH=CH<sub>2</sub> and phenyl-C(=N-)-H.

It is preferred that the protecting group be t-butoxycarbonyl (BOC) and/or benzyloxycarbonyl (CBZ), it is

more preferred that the protecting group be t-butoxycarbonyl. One skilled in the art will recognize suitable methods of introducing a t-butoxycarbonyl or benzyloxycarbonyl protecting group and may additionally consult T.W. Green and P.G.M. Wuts  
5 in "Protective Groups in Organic Chemistry, John Wiley and Sons, 1991 for guidance.

The compounds of the invention may contain geometric or optical isomers as well as tautomers. Thus, the invention includes all tautomers and pure geometric isomers, such as the  
10 *E* and *Z* geometric isomers, as well as mixtures thereof. Further, the invention includes pure enantiomers and diastereomers as well as mixtures thereof, including racemic mixtures. The individual geometric isomers, enantiomers or diastereomers may be prepared or isolated by methods known to  
15 those skilled in the art, including but not limited to chiral chromatography; preparing diastereomers, separating the diastereomers and converting the diastereomers into enantiomers through the use of a chiral resolving agent. A preferred stereochemistry for compounds of formula I is as  
20 follows:



Compounds of the invention with designated stereochemistry can be included in mixtures, including racemic mixtures, with other enantiomers, diastereomers, geometric  
25 isomers or tautomers. In a preferred aspect, compounds of the invention with (S, R, R), (S, S, S), or (S, R, S) stereochemistry are typically present in these mixtures in excess of 50 percent. Preferably, compounds of the invention with designated stereochemistry are present in these mixtures  
30 in excess of 80 percent. More preferably, compounds of the invention with designated stereochemistry are present in these mixtures in excess of 90 percent. Even more preferably,

compounds of the invention with designated stereochemistry are present in these mixtures in excess of 99 percent.

Several of the compounds of formula (I) are amines, and as such form salts when reacted with acids. Pharmaceutically acceptable salts are preferred over the corresponding amines of formula (I) since they produce compounds which are more water soluble, stable and/or more crystalline. Pharmaceutically acceptable salts are any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Pharmaceutically acceptable salts include salts of both inorganic and organic acids. The preferred pharmaceutically acceptable salts include salts of the following acids acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsylic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycollylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, succinic, sulfamic, sulfanilic, sulfonic, sulfuric, tannic, tartaric, teoclic and toluenesulfonic. For other acceptable salts, see *Int. J. Pharm.*, 33, 201-217 (1986) and *J. Pharm. Sci.*, 66(1), 1, (1977).

The invention provides compounds, compositions, kits, and methods for inhibiting beta-secretase enzyme activity and A beta peptide production. Inhibition of beta-secretase enzyme activity halts or reduces the production of A beta from APP

and reduces or eliminates the formation of beta-amyloid deposits in the brain.

#### Methods of the Invention

The compounds of the invention, and pharmaceutically acceptable salts thereof, are useful for treating humans or animals suffering from a condition characterized by a pathological form of beta-amyloid peptide, such as beta-amyloid plaques, and for helping to prevent or delay the onset of such a condition. For example, the compounds are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds of the invention can either be used individually or in combination, as is best for the patient.

As used herein, the term "treating" means that the compounds of the invention can be used in humans with at least a tentative diagnosis of disease. The compounds of the invention will delay or slow the progression of the disease thereby giving the individual a more useful life span.

The term "preventing" means that the compounds of the invention are useful when administered to a patient who has

not been diagnosed as possibly having the disease at the time of administration, but who would normally be expected to develop the disease or be at increased risk for the disease. The compounds of the invention will slow the development of disease symptoms, delay the onset of the disease, or prevent the individual from developing the disease at all. Preventing also includes administration of the compounds of the invention to those individuals thought to be predisposed to the disease due to age, familial history, genetic or chromosomal abnormalities, and/or due to the presence of one or more biological markers for the disease, such as a known genetic mutation of APP or APP cleavage products in brain tissues or fluids.

In treating or preventing the above diseases, the compounds of the invention are administered in a therapeutically effective amount. The therapeutically effective amount will vary depending on the particular compound used and the route of administration, as is known to those skilled in the art.

In treating a patient displaying any of the diagnosed above conditions a physician may administer a compound of the invention immediately and continue administration indefinitely, as needed. In treating patients who are not diagnosed as having Alzheimer's disease, but who are believed to be at substantial risk for Alzheimer's disease, the physician should preferably start treatment when the patient first experiences early pre-Alzheimer's symptoms such as, memory or cognitive problems associated with aging. In addition, there are some patients who may be determined to be at risk for developing Alzheimer's through the detection of a genetic marker such as APOE4 or other biological indicators that are predictive for Alzheimer's disease. In these situations, even though the patient does not have symptoms of the disease, administration of the compounds of the invention may be started before symptoms appear, and treatment may be

continued indefinitely to prevent or delay the onset of the disease.

Dosage Forms and Amounts

5       The compounds of the invention can be administered orally, parenterally, (IV, IM, depo-IM, SQ, and depo SQ), sublingually, intranasally (inhalation), intrathecally, topically, or rectally. Dosage forms known to those of skill in the art are suitable for delivery of the compounds of the  
10 invention.

      Compositions are provided that contain therapeutically effective amounts of the compounds of the invention. The compounds are preferably formulated into suitable pharmaceutical preparations such as tablets, capsules, or  
15 elixirs for oral administration or in sterile solutions or suspensions for parenteral administration. Typically the compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art.

20       About 1 to 500 mg of a compound or mixture of compounds of the invention or a physiologically acceptable salt or ester is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted  
25 pharmaceutical practice. The amount of active substance in those compositions or preparations is such that a suitable dosage in the range indicated is obtained. The compositions are preferably formulated in a unit dosage form, each dosage containing from about 2 to about 100 mg, more preferably about  
30 10 to about 30 mg of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in  
35 association with a suitable pharmaceutical excipient.



To prepare compositions, one or more compounds of the invention are mixed with a suitable pharmaceutically acceptable carrier. Upon mixing or addition of the compound(s), the resulting mixture may be a solution, suspension, emulsion, or the like. Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for lessening or ameliorating at least one symptom of the disease, disorder, or condition treated and may be empirically determined.

Pharmaceutical carriers or vehicles suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. In addition, the active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, or have another action. The compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients.

Where the compounds exhibit insufficient solubility, methods for solubilizing may be used. Such methods are known and include, but are not limited to, using cosolvents such as dimethylsulfoxide (DMSO), using surfactants such as Tween®, and dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as salts or prodrugs may also be used in formulating effective pharmaceutical compositions.

The concentration of the compound is effective for delivery of an amount upon administration that lessens or ameliorates at least one symptom of the disorder for which the

compound is administered. Typically, the compositions are formulated for single dosage administration.

5 The compounds of the invention may be prepared with carriers that protect them against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, microencapsulated delivery systems. The active compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically  
10 useful effect in the absence of undesirable side effects on the patient treated. The therapeutically effective concentration may be determined empirically by testing the compounds in known *in vitro* and *in vivo* model systems for the treated disorder.

15 The compounds and compositions of the invention can be enclosed in multiple or single dose containers. The enclosed compounds and compositions can be provided in kits, for example, including component parts that can be assembled for use. For example, a compound inhibitor in lyophilized form  
20 and a suitable diluent may be provided as separated components for combination prior to use. A kit may include a compound inhibitor and a second therapeutic agent for co-administration. The inhibitor and second therapeutic agent may be provided as separate component parts. A kit may  
25 include a plurality of containers, each container holding one or more unit dose of the compound of the invention. The containers are preferably adapted for the desired mode of administration, including, but not limited to tablets, gel capsules, sustained-release capsules, and the like for oral  
30 administration; depot products, pre-filled syringes, ampoules, vials, and the like for parenteral administration; and patches, medipads, creams, and the like for topical administration.

The concentration of active compound in the drug  
35 composition will depend on absorption, inactivation, and

excretion rates of the active compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

The active ingredient may be administered at once, or may  
5 be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro*  
10 test data. It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the  
15 professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

20 If oral administration is desired, the compound should be provided in a composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an enteric coating that maintains its integrity in the stomach and releases the active compound in  
25 the intestine. The composition may also be formulated in combination with an antacid or other such ingredient.

Oral compositions will generally include an inert diluent or an edible carrier and may be compressed into tablets or enclosed in gelatin capsules. For the purpose of oral  
30 therapeutic administration, the active compound or compounds can be incorporated with excipients and used in the form of tablets, capsules, or troches. Pharmaceutically compatible binding agents and adjuvant materials can be included as part of the composition.

The tablets, pills, capsules, troches, and the like can contain any of the following ingredients or compounds of a similar nature: a binder such as, but not limited to, gum tragacanth, acacia, corn starch, or gelatin; an excipient such as microcrystalline cellulose, starch, or lactose; a disintegrating agent such as, but not limited to, alginic acid and corn starch; a lubricant such as, but not limited to, magnesium stearate; a gildant, such as, but not limited to, colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; and a flavoring agent such as peppermint, methyl salicylate, or fruit flavoring.

When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials, which modify the physical form of the dosage unit, for example, coatings of sugar and other enteric agents. The compounds can also be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings, and flavors.

The active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action.

Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent such as water for injection, saline solution, fixed oil, a naturally occurring vegetable oil such as sesame oil, coconut oil, peanut oil, cottonseed oil, and the like, or a synthetic fatty vehicle such as ethyl oleate, and the like, polyethylene glycol, glycerine, propylene glycol, or other synthetic solvent; antimicrobial agents such as benzyl alcohol and methyl parabens; antioxidants such as ascorbic acid and sodium bisulfite; chelating agents such as ethylenediaminetetraacetic

acid (EDTA); buffers such as acetates, citrates, and phosphates; and agents for the adjustment of tonicity such as sodium chloride and dextrose. Parenteral preparations can be enclosed in ampoules, disposable syringes, or multiple dose  
5 vials made of glass, plastic, or other suitable material. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

Where administered intravenously, suitable carriers include physiological saline, phosphate buffered saline (PBS),  
10 and solutions containing thickening and solubilizing agents such as glucose, polyethylene glycol, polypropyleneglycol, and mixtures thereof. Liposomal suspensions including tissue-targeted liposomes may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to  
15 methods known for example, as described in U.S. Patent No. 4,522,811.

The active compounds may be prepared with carriers that protect the compound against rapid elimination from the body, such as time-release formulations or coatings. Such carriers  
20 include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers such as collagen, ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid, and the like. Methods for  
25 preparation of such formulations are known to those skilled in the art.

The compounds of the invention can be administered orally, parenterally (IV, IM, depo-IM, SQ, and depo-SQ), sublingually, intranasally (inhalation), intrathecally,  
30 topically, or rectally. Dosage forms known to those skilled in the art are suitable for delivery of the compounds of the invention.

Compounds of the invention may be administered enterally or parenterally. When administered orally, compounds of the  
35 invention can be administered in usual dosage forms for oral

administration as is well known to those skilled in the art. These dosage forms include the usual solid unit dosage forms of tablets and capsules as well as liquid dosage forms such as solutions, suspensions, and elixirs. When the solid dosage forms are used, it is preferred that they be of the sustained release type so that the compounds of the invention need to be administered only once or twice daily.

The oral dosage forms are administered to the patient 1, 2, 3, or 4 times daily. It is preferred that the compounds of the invention be administered either three or fewer times, more preferably once or twice daily. Hence, it is preferred that the compounds of the invention be administered in oral dosage form. It is preferred that whatever oral dosage form is used, that it be designed so as to protect the compounds of the invention from the acidic environment of the stomach. Enteric coated tablets are well known to those skilled in the art. In addition, capsules filled with small spheres each coated to protect from the acidic stomach, are also well known to those skilled in the art.

When administered orally, an administered amount therapeutically effective to inhibit beta-secretase activity, to inhibit A beta production, to inhibit A beta deposition, or to treat or prevent AD is from about 0.1 mg/day to about 1,000 mg/day. It is preferred that the oral dosage is from about 1 mg/day to about 100 mg/day. It is more preferred that the oral dosage is from about 5 mg/day to about 50 mg/day. It is understood that while a patient may be started at one dose, that dose may be varied over time as the patient's condition changes.

Compounds of the invention may also be advantageously delivered in a nano crystal dispersion formulation. Preparation of such formulations is described, for example, in U.S. Patent 5,145,684. Nano crystalline dispersions of HIV protease inhibitors and their method of use are described in

U.S. Patent No. 6,045,829. The nano crystalline formulations typically afford greater bioavailability of drug compounds.

The compounds of the invention can be administered parenterally, for example, by IV, IM, depo-IM, SC, or depo-SC.

5 When administered parenterally, a therapeutically effective amount of about 0.5 to about 100 mg/day, preferably from about 5 to about 50 mg daily should be delivered. When a depot formulation is used for injection once a month or once every two weeks, the dose should be about 0.5 mg/day to about 50  
10 mg/day, or a monthly dose of from about 15 mg to about 1,500 mg. In part because of the forgetfulness of the patients with Alzheimer's disease, it is preferred that the parenteral dosage form be a depo formulation.

The compounds of the invention can be administered  
15 sublingually. When given sublingually, the compounds of the invention should be given one to four times daily in the amounts described above for IM administration.

The compounds of the invention can be administered intranasally. When given by this route, the appropriate  
20 dosage forms are a nasal spray or dry powder, as is known to those skilled in the art. The dosage of the compounds of the invention for intranasal administration is the amount described above for IM administration.

The compounds of the invention can be administered  
25 intrathecally. When given by this route the appropriate dosage form can be a parenteral dosage form as is known to those skilled in the art. The dosage of the compounds of the invention for intrathecal administration is the amount described above for IM administration.

30 The compounds of the invention can be administered topically. When given by this route, the appropriate dosage form is a cream, ointment, or patch. Because of the amount of the compounds of the invention to be administered, the patch is preferred. When administered topically, the dosage is from  
35 about 0.5 mg/day to about 200 mg/day. Because the amount that

can be delivered by a patch is limited, two or more patches may be used. The number and size of the patch is not important, what is important is that a therapeutically effective amount of the compounds of the invention be  
5 delivered as is known to those skilled in the art. The compounds of the invention can be administered rectally by suppository as is known to those skilled in the art. When administered by suppository, the therapeutically effective amount is from about 0.5 mg to about 500 mg.

10 The compounds of the invention can be administered by implants as is known to those skilled in the art. When administering a compound of the invention by implant, the therapeutically effective amount is the amount described above for depot administration.

15 Given a particular compound of the invention and a desired dosage form, one skilled in the art would know how to prepare and administer the appropriate dosage form.

The compounds of the invention are used in the same manner, by the same routes of administration, using the same  
20 pharmaceutical dosage forms, and at the same dosing schedule as described above, for preventing disease or treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating or preventing  
25 Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias,  
30 including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type of Alzheimer's disease.



The compounds of the invention can be used in combination, with each other or with other therapeutic agents or approaches used to treat or prevent the conditions listed above. Such agents or approaches include: acetylcholine  
5 esterase inhibitors such as tacrine (tetrahydroaminoacridine, marketed as COGNEX®), donepezil hydrochloride, (marketed as Aricept® and rivastigmine (marketed as Exelon®); gamma-secretase inhibitors; anti-inflammatory agents such as cyclooxygenase II inhibitors; anti-oxidants such as Vitamin E  
10 and ginkgolides; immunological approaches, such as, for example, immunization with A beta peptide or administration of anti-A beta peptide antibodies; statins; and direct or indirect neurotropic agents such as Cerebrolysin®, AIT-082 (Emilieu, 2000, Arch. Neurol. 57:454), and other neurotropic  
15 agents of the future.

In addition, the compounds of formula (I) can also be used with inhibitors of P-glycoprotein (P-gp). P-gp inhibitors and the use of such compounds are known to those skilled in the art. See for example, *Cancer Research*, 53,  
20 4595-4602 (1993), *Clin. Cancer Res.*, 2, 7-12 (1996), *Cancer Research*, 56, 4171-4179 (1996), International Publications WO99/64001 and WO01/10387. The important thing is that the blood level of the P-gp inhibitor be such that it exerts its effect in inhibiting P-gp from decreasing brain blood levels  
25 of the compounds of formula (A). To that end the P-gp inhibitor and the compounds of formula (A) can be administered at the same time, by the same or different route of administration, or at different times. The important thing is not the time of administration but having an effective blood  
30 level of the P-gp inhibitor.

Suitable P-gp inhibitors include cyclosporin A, verapamil, tamoxifen, quinidine, Vitamin E-TGPS, ritonavir, megestrol acetate, progesterone, rapamycin, 10,11-methanodibenzosuberane, phenothiazines, acridine derivatives  
35 such as GF120918, FK506, VX-710, LY335979, PSC-833, GF-102,918

and other steroids. It is to be understood that additional agents will be found that have the same function and therefore achieve the same outcome; such compounds are also considered to be useful.

5       The P-gp inhibitors can be administered orally, parenterally, (IV, IM, IM-depo, SQ, SQ-depo), topically, sublingually, rectally, intranasally, intrathecally and by implant.

10       The therapeutically effective amount of the P-gp inhibitors is from about 0.1 to about 300 mg/kg/day, preferably about 0.1 to about 150 mg/kg daily. It is understood that while a patient may be started on one dose, that dose may have to be varied over time as the patient's condition changes.

15       When administered orally, the P-gp inhibitors can be administered in usual dosage forms for oral administration as is known to those skilled in the art. These dosage forms include the usual solid unit dosage forms of tablets and capsules as well as liquid dosage forms such as solutions, 20       suspensions and elixirs. When the solid dosage forms are used, it is preferred that they be of the sustained release type so that the P-gp inhibitors need to be administered only once or twice daily. The oral dosage forms are administered to the patient one thru four times daily. It is preferred 25       that the P-gp inhibitors be administered either three or fewer times a day, more preferably once or twice daily. Hence, it is preferred that the P-gp inhibitors be administered in solid dosage form and further it is preferred that the solid dosage form be a sustained release form which permits once or twice 30       daily dosing. It is preferred that what ever dosage form is used, that it be designed so as to protect the P-gp inhibitors from the acidic environment of the stomach. Enteric coated tablets are well known to those skilled in the art. In addition, capsules filled with small spheres each coated to

protect from the acidic stomach, are also well known to those skilled in the art.

In addition, the P-gp inhibitors can be administered parenterally. When administered parenterally they can be administered IV, IM, depo-IM, SQ or depo-SQ.

The P-gp inhibitors can be given sublingually. When given sublingually, the P-gp inhibitors should be given one thru four times daily in the same amount as for IM administration.

The P-gp inhibitors can be given intranasally. When given by this route of administration, the appropriate dosage forms are a nasal spray or dry powder as is known to those skilled in the art. The dosage of the P-gp inhibitors for intranasal administration is the same as for IM administration.

The P-gp inhibitors can be given intrathecally. When given by this route of administration the appropriate dosage form can be a parenteral dosage form as is known to those skilled in the art.

The P-gp inhibitors can be given topically. When given by this route of administration, the appropriate dosage form is a cream, ointment or patch. Because of the amount of the P-gp inhibitors needed to be administered the patch is preferred. However, the amount that can be delivered by a patch is limited. Therefore, two or more patches may be required. The number and size of the patch is not important, what is important is that a therapeutically effective amount of the P-gp inhibitors be delivered as is known to those skilled in the art.

The P-gp inhibitors can be administered rectally by suppository as is known to those skilled in the art.

The P-gp inhibitors can be administered by implants as is known to those skilled in the art.

There is nothing novel about the route of administration nor the dosage forms for administering the P-gp inhibitors.

Given a particular P-gp inhibitor, and a desired dosage form, one skilled in the art would know how to prepare the appropriate dosage form for the P-gp inhibitor.

It should be apparent to one skilled in the art that the exact dosage and frequency of administration will depend on the particular compounds of the invention administered, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, and other medication the individual may be taking as is well known to administering physicians who are skilled in this art.

#### Inhibition of APP Cleavage

The compounds of the invention inhibit cleavage of APP between Met595 and Asp596 numbered for the APP695 isoform, or a mutant thereof, or at a corresponding site of a different isoform, such as APP751 or APP770, or a mutant thereof (sometimes referred to as the "beta secretase site"). While not wishing to be bound by a particular theory, inhibition of beta-secretase activity is thought to inhibit production of beta amyloid peptide (A beta). Inhibitory activity is demonstrated in one of a variety of inhibition assays, whereby cleavage of an APP substrate in the presence of a beta-secretase enzyme is analyzed in the presence of the inhibitory compound, under conditions normally sufficient to result in cleavage at the beta-secretase cleavage site. Reduction of APP cleavage at the beta-secretase cleavage site compared with an untreated or inactive control is correlated with inhibitory activity. Assay systems that can be used to demonstrate efficacy of the compound inhibitors of the invention are known. Representative assay systems are described, for example, in U.S. Patents No. 5,942,400, 5,744,346, as well as in the Examples below.

The enzymatic activity of beta-secretase and the production of A beta can be analyzed *in vitro* or *in vivo*,

using natural, mutated, and/or synthetic APP substrates, natural, mutated, and/or synthetic enzyme, and the test compound. The analysis may involve primary or secondary cells expressing native, mutant, and/or synthetic APP and enzyme, animal models expressing native APP and enzyme, or may utilize transgenic animal models expressing the substrate and enzyme. Detection of enzymatic activity can be by analysis of one or more of the cleavage products, for example, by immunoassay, fluorometric or chromogenic assay, HPLC, or other means of detection. Inhibitory compounds are determined as those having the ability to decrease the amount of beta-secretase cleavage product produced in comparison to a control, where beta-secretase mediated cleavage in the reaction system is observed and measured in the absence of inhibitory compounds.

#### Beta-Secretase

Various forms of beta-secretase enzyme are known, and are available and useful for assay of enzyme activity and inhibition of enzyme activity. These include native, recombinant, and synthetic forms of the enzyme. Human beta-secretase is known as Beta Site APP Cleaving Enzyme (BACE), Asp2, and memapsin 2, and has been characterized, for example, in U.S. Patent No. 5,744,346 and published PCT patent applications WO98/22597, WO00/03819, WO01/23533, and WO00/17369, as well as in literature publications (Hussain et al., 1999, *Mol. Cell. Neurosci.* 14:419-427; Vassar et al., 1999, *Science* 286:735-741; Yan et al., 1999, *Nature* 402:533-537; Sinha et al., 1999, *Nature* 40:537-540; and Lin et al., 2000, *PNAS USA* 97:1456-1460). Synthetic forms of the enzyme have also been described (WO98/22597 and WO00/17369). Beta-secretase can be extracted and purified from human brain tissue and can be produced in cells, for example mammalian cells expressing recombinant enzyme.

Preferred compounds are effective to inhibit 50% of beta-secretase enzymatic activity at a concentration of less than

50 micromolar, preferably at a concentration of 10 micromolar or less, more preferably 1 micromolar or less, and most preferably 10 nanomolar or less.

#### 5           APP Substrate

Assays that demonstrate inhibition of beta-secretase-mediated cleavage of APP can utilize any of the known forms of APP, including the 695 amino acid "normal" isotype described by Kang et al., 1987, *Nature* 325:733-6, the 770 amino acid isotype described by Kitaguchi et. al., 1981, *Nature* 331:530-532, and variants such as the Swedish Mutation (KM670-1NL) (APP-SW), the London Mutation (V7176F), and others. See, for example, U.S. Patent No. 5,766,846 and also Hardy, 1992, *Nature Genet.* 1:233-234, for a review of known variant mutations. Additional useful substrates include the dibasic amino acid modification, APP-KK disclosed, for example, in WO 00/17369, fragments of APP, and synthetic peptides containing the beta-secretase cleavage site, wild type (WT) or mutated form, e.g., SW, as described, for example, in U.S. Patent No 20 5,942,400 and WO00/03819.

The APP substrate contains the beta-secretase cleavage site of APP (KM-DA or NL-DA) for example, a complete APP peptide or variant, an APP fragment, a recombinant or synthetic APP, or a fusion peptide. Preferably, the fusion peptide includes the beta-secretase cleavage site fused to a peptide having a moiety useful for enzymatic assay, for example, having isolation and/or detection properties. A useful moiety may be an antigenic epitope for antibody binding, a label or other detection moiety, a binding substrate, and the like. 25 30

#### Antibodies

Products characteristic of APP cleavage can be measured by immunoassay using various antibodies, as described, for example, in Pirttila et al., 1999, *Neuro. Lett.* 249:21-4, and 35

in U.S. Patent No. 5,612,486. Useful antibodies to detect A beta include, for example, the monoclonal antibody 6E10 (Senetek, St. Louis, MO) that specifically recognizes an epitope on amino acids 1-16 of the A beta peptide; antibodies 5 162 and 164 (New York State Institute for Basic Research, Staten Island, NY) that are specific for human A beta 1-40 and 1-42, respectively; and antibodies that recognize the junction region of beta-amyloid peptide, the site between residues 16 and 17, as described in U.S. Patent No. 5,593,846. Antibodies 10 raised against a synthetic peptide of residues 591 to 596 of APP and SW192 antibody raised against 590-596 of the Swedish mutation are also useful in immunoassay of APP and its cleavage products, as described in U.S. Patent Nos. 5,604,102 and 5,721,130.

#### Assay Systems

Assays for determining APP cleavage at the beta-secretase cleavage site are well known in the art. Exemplary assays, are described, for example, in U.S. Patent Nos. 5,744,346 and 20 5,942,400, and described in the Examples below.

#### Cell Free Assays

Exemplary assays that can be used to demonstrate the inhibitory activity of the compounds of the invention are 25 described, for example, in WO00/17369, WO 00/03819, and U.S. Patents No. 5,942,400 and 5,744,346. Such assays can be performed in cell-free incubations or in cellular incubations using cells expressing a beta-secretase and an APP substrate having a beta-secretase cleavage site.

30 An APP substrate containing the beta-secretase cleavage site of APP, for example, a complete APP or variant, an APP fragment, or a recombinant or synthetic APP substrate containing the amino acid sequence: KM-DA or NL-DA, is incubated in the presence of beta-secretase enzyme, a 35 fragment thereof, or a synthetic or recombinant polypeptide

variant having beta-secretase activity and effective to cleave the beta-secretase cleavage site of APP, under incubation conditions suitable for the cleavage activity of the enzyme. Suitable substrates optionally include derivatives that may be fusion proteins or peptides that contain the substrate peptide and a modification useful to facilitate the purification or detection of the peptide or its beta-secretase cleavage products. Useful modifications include the insertion of a known antigenic epitope for antibody binding; the linking of a label or detectable moiety, the linking of a binding substrate, and the like.

Suitable incubation conditions for a cell-free *in vitro* assay include, for example: approximately 200 nanomolar to 10 micromolar substrate, approximately 10 to 200 picomolar enzyme, and approximately 0.1 nanomolar to 10 micromolar inhibitor compound, in aqueous solution, at an approximate pH of 4 -7, at approximately 37 degrees C, for a time period of approximately 10 minutes to 3 hours. These incubation conditions are exemplary only, and can be varied as required for the particular assay components and/or desired measurement system. Optimization of the incubation conditions for the particular assay components should account for the specific beta-secretase enzyme used and its pH optimum, any additional enzymes and/or markers that might be used in the assay, and the like. Such optimization is routine and will not require undue experimentation.

One useful assay utilizes a fusion peptide having maltose binding protein (MBP) fused to the C-terminal 125 amino acids of APP-SW. The MBP portion is captured on an assay substrate by anti-MBP capture antibody. Incubation of the captured fusion protein in the presence of beta-secretase results in cleavage of the substrate at the beta-secretase cleavage site. Analysis of the cleavage activity can be, for example, by immunoassay of cleavage products. One such immunoassay detects a unique epitope exposed at the carboxy terminus of



the cleaved fusion protein, for example, using the antibody SW192. This assay is described, for example, in U.S. Patent No 5,942,400.

## 5 Cellular Assay

Numerous cell-based assays can be used to analyze beta-secretase activity and/or processing of APP to release A beta. Contact of an APP substrate with a beta-secretase enzyme within the cell and in the presence or absence of a compound  
10 inhibitor of the invention can be used to demonstrate beta-secretase inhibitory activity of the compound. Preferably, assay in the presence of a useful inhibitory compound provides at least about 30%, most preferably at least about 50% inhibition of the enzymatic activity, as compared with a non-  
15 inhibited control.

In one embodiment, cells that naturally express beta-secretase are used. Alternatively, cells are modified to express a recombinant beta-secretase or synthetic variant enzyme as discussed above. The APP substrate may be added to  
20 the culture medium and is preferably expressed in the cells. Cells that naturally express APP, variant or mutant forms of APP, or cells transformed to express an isoform of APP, mutant or variant APP, recombinant or synthetic APP, APP fragment, or synthetic APP peptide or fusion protein containing the beta-  
25 secretase APP cleavage site can be used, provided that the expressed APP is permitted to contact the enzyme and enzymatic cleavage activity can be analyzed.

Human cell lines that normally process A beta from APP provide a useful means to assay inhibitory activities of the  
30 compounds of the invention. Production and release of A beta and/or other cleavage products into the culture medium can be measured, for example by immunoassay, such as Western blot or enzyme-linked immunoassay (EIA) such as by ELISA.

Cells expressing an APP substrate and an active beta-  
35 secretase can be incubated in the presence of a compound

inhibitor to demonstrate inhibition of enzymatic activity as compared with a control. Activity of beta-secretase can be measured by analysis of one or more cleavage products of the APP substrate. For example, inhibition of beta-secretase activity against the substrate APP would be expected to decrease release of specific beta-secretase induced APP cleavage products such as A beta.

Although both neural and non-neural cells process and release A beta, levels of endogenous beta-secretase activity are low and often difficult to detect by EIA. The use of cell types known to have enhanced beta-secretase activity, enhanced processing of APP to A beta, and/or enhanced production of A beta are therefore preferred. For example, transfection of cells with the Swedish Mutant form of APP (APP-SW); with APP-KK; or with APP-SW-KK provides cells having enhanced beta-secretase activity and producing amounts of A beta that can be readily measured.

In such assays, for example, the cells expressing APP and beta-secretase are incubated in a culture medium under conditions suitable for beta-secretase enzymatic activity at its cleavage site on the APP substrate. On exposure of the cells to the compound inhibitor, the amount of A beta released into the medium and/or the amount of CTF99 fragments of APP in the cell lysates is reduced as compared with the control. The cleavage products of APP can be analyzed, for example, by immune reactions with specific antibodies, as discussed above.

Preferred cells for analysis of beta-secretase activity include primary human neuronal cells, primary transgenic animal neuronal cells where the transgene is APP, and other cells such as those of a stable 293 cell line expressing APP, for example, APP-SW.

#### **In vivo Assays: Animal Models**

Various animal models can be used to analyze beta-secretase activity and /or processing of APP to release A

beta, as described above. For example, transgenic animals expressing APP substrate and beta-secretase enzyme can be used to demonstrate inhibitory activity of the compounds of the invention. Certain transgenic animal models have been described, for example, in U.S. Patent Nos.: 5,877,399; 5,612,486; 5,387,742; 5,720,936; 5,850,003; 5,877,015,, and 5,811,633, and in Ganes et al., 1995, Nature 373:523. Preferred are animals that exhibit characteristics associated with the pathophysiology of AD. Administration of the compound inhibitors of the invention to the transgenic mice described herein provides an alternative method for demonstrating the inhibitory activity of the compounds. Administration of the compounds in a pharmaceutically effective carrier and via an administrative route that reaches the target tissue in an appropriate therapeutic amount is also preferred.

Inhibition of beta-secretase mediated cleavage of APP at the beta-secretase cleavage site and of A beta release can be analyzed in these animals by measure of cleavage fragments in the animal's body fluids such as cerebral fluid or tissues. Analysis of brain tissues for A beta deposits or plaques is preferred.

On contacting an APP substrate with a beta-secretase enzyme in the presence of an inhibitory compound of the invention and under conditions sufficient to permit enzymatic mediated cleavage of APP and/or release of A beta from the substrate, the compounds of the invention are effective to reduce beta-secretase-mediated cleavage of APP at the beta-secretase cleavage site and/or effective to reduce released amounts of A beta. Where such contacting is the administration of the inhibitory compounds of the invention to an animal model, for example, as described above, the compounds are effective to reduce A beta deposition in brain tissues of the animal, and to reduce the number and/or size of beta amyloid plaques. Where such administration is to a human

subject, the compounds are effective to inhibit or slow the progression of disease characterized by enhanced amounts of A beta, to slow the progression of AD in the, and/or to prevent onset or development of AD in a patient at risk for the disease.

Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are hereby incorporated by reference for all purposes.

### Definitions

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims:

It should be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to a composition containing "a compound" includes a mixture of two or more compounds. It should also be noted that the term "or" is generally employed in its sense including "and/or" unless the content clearly dictates otherwise.

The symbol "-" in general represents a bond between two atoms in the chain. Thus  $\text{CH}_3\text{-O-CH}_2\text{-CH(R}_1\text{)-CH}_3$  represents a 2-substituted-1-methoxypropane compound. In addition, the symbol "-" represents the point of attachment of the substituent to a compound. Thus for example  $\text{aryl(C}_1\text{-C}_6\text{)alkyl-}$  indicates an alkylaryl group, such as benzyl, attached to the compound at the alkyl moiety.

Where multiple substituents are indicated as being attached to a structure, it is to be understood that the substituents can be the same or different. Thus for example "R<sub>m</sub> optionally substituted with 1, 2 or 3 R<sub>q</sub> groups" indicates

that  $R_m$  is substituted with 1, 2, or 3  $R_q$  groups where the  $R_q$  groups can be the same or different.

APP, amyloid precursor protein, is defined as any APP polypeptide, including APP variants, mutations, and isoforms, for example, as disclosed in U.S. Patent No. 5,766,846.

A beta, amyloid beta peptide, is defined as any peptide resulting from beta-secretase mediated cleavage of APP, including peptides of 39, 40, 41, 42, and 43 amino acids, and extending from the beta-secretase cleavage site to amino acids 39, 40, 41, 42, or 43.

Beta-secretase (BACE1, Asp2, Memapsin 2) is an aspartyl protease that mediates cleavage of APP at the amino-terminal edge of A beta. Human beta-secretase is described, for example, in WO00/17369.

Pharmaceutically acceptable refers to those properties and/or substances that are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

A therapeutically effective amount is defined as an amount effective to reduce or lessen at least one symptom of the disease being treated or to reduce or delay onset of one or more clinical markers or symptoms of the disease.

By "alkyl" and " $C_1$ - $C_6$  alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl. It is understood that in cases where an alkyl chain of a substituent (e.g. of an alkyl, alkoxy or alkenyl group) is shorter or longer than 6 carbons, it will be so indicated in the second "C" as, for example, " $C_1$ - $C_{10}$ " indicates a maximum of 10 carbons.

By "alkoxy" and " $C_1$ - $C_6$  alkoxy" in the present invention is meant straight or branched chain alkyl groups having 1-6

carbon atoms, attached through at least one divalent oxygen atom, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, isopentoxy, neopentoxy, hexoxy, and 3-methylpentoxy.

5 By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

"Alkenyl" and "C<sub>2</sub>-C<sub>6</sub> alkenyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and from one to three double bonds and includes, for example, ethenyl, propenyl, 1-but-3-enyl, 1-pent-3-enyl, 1-hex-5-enyl and the like.

"Alkynyl" and "C<sub>2</sub>-C<sub>6</sub> alkynyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and one or two triple bonds and includes ethynyl, propynyl, butynyl, pentyn-2-yl and the like.

As used herein, the term "cycloalkyl" refers to saturated carbocyclic radicals having three to twelve carbon atoms. The cycloalkyl can be monocyclic, or a polycyclic fused system. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The cycloalkyl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such cycloalkyl groups may be optionally substituted with C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

By "aryl" is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl), which is optionally mono-, di-, or trisubstituted. Preferred aryl groups of the present invention are phenyl, 1-naphthyl, 2-naphthyl, indanyl, indenyl, dihydronaphthyl, tetralinyl or

6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl. The aryl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such aryl groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, -COOH, -C(=O)O(C<sub>1</sub>-C<sub>6</sub> alkyl), -C(=O)NH<sub>2</sub>, -C(=O)N(mono- or di-C<sub>1</sub>-C<sub>6</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -O-C(=O)(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)NH<sub>2</sub>, -NH-C(=O)N(mono- or di-C<sub>1</sub>-C<sub>6</sub> alkyl), -NH(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-NH<sub>2</sub> or -NH(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-N-(mono- or di-C<sub>1</sub>-C<sub>6</sub> alkyl).

By "heteroaryl" is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heteroaryl groups of the present invention include pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indolizinyl, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, isochromanyl, chromanyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranlyl, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranlyl, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl,

benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl,  
 benzopyranyl, benzothiopyranyl, coumarinyl, isocoumarinyl,  
 chromonyl, chromanonyl, pyridinyl-N-oxide,  
 tetrahydroquinolinyl, dihydroquinolinyl, dihydroquinolinonyl,  
 5 dihydroisoquinolinonyl, dihydrocoumarinyl,  
 dihydroisocoumarinyl, isoindolinonyl, benzodioxanyl,  
 benzoxazolinonyl, pyrrolyl N-oxide,, pyrimidinyl N-oxide,  
 pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide,  
 indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide,  
 10 quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-  
 oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-  
 oxide, thiazolyl N-oxide, indoliziny N-oxide, indazolyl N-  
 oxide, benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl  
 N-oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl  
 15 N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide,  
 benzothiopyranyl S,S-dioxide. The heteroaryl groups herein  
 are unsubstituted or, as specified, substituted in one or more  
 substitutable positions with various groups. For example,  
 such heteroaryl groups may be optionally substituted with C<sub>1</sub>-C<sub>6</sub>  
 20 alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino,  
 mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-  
 C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-  
 C<sub>6</sub>)alkyl, -COOH, -C(=O)O(C<sub>1</sub>-C<sub>6</sub> alkyl), -C(=O)NH<sub>2</sub>, -C(=O)N(mono-  
 25 or di-C<sub>1</sub>-C<sub>6</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), -O-  
 C(=O)(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)-  
 C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)-SO<sub>2</sub>-  
 (C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)NH<sub>2</sub>, -NH-C(=O)N(mono- or di-C<sub>1</sub>-C<sub>6</sub>  
 alkyl), -NH(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-NH<sub>2</sub> or -NH(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-N-  
 30 (mono- or di-C<sub>1</sub>-C<sub>6</sub> alkyl).

By "heterocycle", "heterocycloalkyl" or "heterocyclyl"  
 is meant one or more carbocyclic ring systems of 3-, 4-, 5-,  
 6-, or 7-membered rings which includes fused ring systems of  
 9-11 atoms containing at least one and up to four heteroatoms  
 35 selected from nitrogen, oxygen, or sulfur. Preferred



heterocycles of the present invention include morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, 5 tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide, oxazolidinonyl, dihydropyrazolyl, dihydroisoindolyl, dihydropyrrolyl, dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, azepanyl, 10 diazepanyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide and homothiomorpholinyl S-oxide. The heterocycle groups herein maybe unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such heterocycle groups may be 15 optionally substituted with C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or =O.

20 All patents and publications referred to herein are hereby incorporated by reference for all purposes.

Structures were named using Name Pro IUPAC Naming Software, version 5.09, available from Advanced Chemical Development, Inc., 90 Adelaide Street West, Toronto, Ontario, 25 M5H 3V9, Canada, or using Chemdraw Ultra version 6.0.2, which is available through Cambridgesoft, 100 Cambridge Park Drive, Cambridge, MA 02140, or were derived therefrom.

The present invention may be better understood with reference to the following examples. These examples are 30 intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

BIOLOGY EXAMPLESExample AEnzyme Inhibition Assay

The compounds of the invention are analyzed for inhibitory activity by use of the MBP-C125 assay. This assay determines the relative inhibition of beta-secretase cleavage of a model APP substrate, MBP-C125SW, by the compounds assayed as compared with an untreated control. A detailed description of the assay parameters can be found, for example, in U.S. Patent No. 5,942,400. Briefly, the substrate is a fusion peptide formed of maltose binding protein (MBP) and the carboxy terminal 125 amino acids of APP-SW, the Swedish mutation. The beta-secretase enzyme is derived from human brain tissue as described in Sinha et al, 1999, *Nature* 40:537-540) or recombinantly produced as the full-length enzyme (amino acids 1-501), and can be prepared, for example, from 293 cells expressing the recombinant cDNA, as described in WO00/47618.

Inhibition of the enzyme is analyzed, for example, by immunoassay of the enzyme's cleavage products. One exemplary ELISA uses an anti-MBP capture antibody that is deposited on precoated and blocked 96-well high binding plates, followed by incubation with diluted enzyme reaction supernatant, incubation with a specific reporter antibody, for example, biotinylated anti-SW192 reporter antibody, and further incubation with streptavidin/alkaline phosphatase. In the assay, cleavage of the intact MBP-C125SW fusion protein results in the generation of a truncated amino-terminal fragment, exposing a new SW-192 antibody-positive epitope at the carboxy terminus. Detection is effected by a fluorescent substrate signal on cleavage by the phosphatase. ELISA only detects cleavage following Leu 596 at the substrate's APP-SW 751 mutation site.

**Specific Assay Procedure:**

Compounds are diluted in a 1:1 dilution series to a six-point concentration curve (two wells per concentration) in one 96-plate row per compound tested. Each of the test compounds is prepared in DMSO to make up a 10 millimolar stock solution. The stock solution is serially diluted in DMSO to obtain a final compound concentration of 200 micromolar at the high point of a 6-point dilution curve. Ten (10) microliters of each dilution is added to each of two wells on row C of a corresponding V-bottom plate to which 190 microliters of 52 millimolar NaOAc, 7.9% DMSO, pH 4.5 are pre-added. The NaOAc diluted compound plate is spun down to pellet precipitant and 20 microliters/well is transferred to a corresponding flat-bottom plate to which 30 microliters of ice-cold enzyme-substrate mixture (2.5 microliters MBP-C125SW substrate, 0.03 microliters enzyme and 24.5 microliters ice cold 0.09% TX100 per 30 microliters) is added. The final reaction mixture of 200 micromolar compound at the highest curve point is in 5% DMSO, 20 millimolar NaOAc, 0.06% TX100, at pH 4.5.

Warming the plates to 37 degrees C starts the enzyme reaction. After 90 minutes at 37 degrees C, 200 microliters/well cold specimen diluent is added to stop the reaction and 20 microliters/well was transferred to a corresponding anti-MBP antibody coated ELISA plate for capture, containing 80 microliters/well specimen diluent. This reaction is incubated overnight at 4 degrees C and the ELISA is developed the next day after a 2 hour incubation with anti-192SW antibody, followed by Streptavidin-AP conjugate and fluorescent substrate. The signal is read on a fluorescent plate reader.

Relative compound inhibition potency is determined by calculating the concentration of compound that showed a fifty percent reduction in detected signal ( $IC_{50}$ ) compared to the enzyme reaction signal in the control wells with no added

compound. In this assay, preferred compounds of the invention exhibit an IC<sub>50</sub> of less than 50 micromolar.

#### Example B

#### 5      Cell Free Inhibition Assay Utilizing a Synthetic APP Substrate

A synthetic APP substrate that can be cleaved by beta-secretase and having N-terminal biotin and made fluorescent by the covalent attachment of Oregon green at the Cys residue is  
10      used to assay beta-secretase activity in the presence or absence of the inhibitory compounds of the invention. Useful substrates include the following:

Biotin-SEVNL-DAEFRC [oregon green] KK      [SEQ ID NO:  
15      1]  
Biotin-SEVKM-DAEFRC [oregon green] KK      [SEQ ID NO:  
2]  
Biotin-GLNIKTEEISEISY-EVEFRC [oregon green] KK [SEQ ID NO:  
3]  
20      Biotin-ADRGLTTRPGSGLTNIKTEEISEVNL-DAEFRC [oregon green] KK  
[SEQ ID NO:4]  
Biotin-FVNQHLCoxGSHLVEALY-LVCoxGERGFFYTPKAC [oregon  
green] KK      [SEQ ID NO: 5]

The enzyme (0.1 nanomolar) and test compounds (0.001 -  
25      100 micromolar) are incubated in pre-blocked, low affinity, black plates (384 well) at 37 degrees for 30 minutes. The reaction is initiated by addition of 150 millimolar substrate to a final volume of 30 microliter per well. The final assay conditions are: 0.001 - 100 micromolar compound inhibitor;  
30      0.1 molar sodium acetate (pH 4.5); 150 nanomolar substrate; 0.1 nanomolar soluble beta-secretase; 0.001% Tween 20, and 2% DMSO. The assay mixture is incubated for 3 hours at 37 degrees C, and the reaction is terminated by the addition of a saturating concentration of immunopure streptavidin. After  
35      incubation with streptavidin at room temperature for 15

minutes, fluorescence polarization is measured, for example, using a LJL Acquest (Ex485 nm/ Em530 nm). The activity of the beta-secretase enzyme is detected by changes in the fluorescence polarization that occur when the substrate is cleaved by the enzyme. Incubation in the presence or absence of compound inhibitor demonstrates specific inhibition of beta-secretase enzymatic cleavage of its synthetic APP substrate. In this assay, preferred compounds of the invention exhibit an IC<sub>50</sub> of less than 50 micromolar.

#### Example C

##### Beta-Secretase Inhibition: P26-P4'SW Assay

Synthetic substrates containing the beta-secretase cleavage site of APP are used to assay beta-secretase activity, using the methods described, for example, in published PCT application WO00/47618. The P26-P4'SW substrate is a peptide of the sequence:

(biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNLDAEF [SEQ ID NO: 6]

The P26-P1 standard has the sequence:

(biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNL [SEQ ID NO: 7].

Briefly, the biotin-coupled synthetic substrates are incubated at a concentration of from about 0 to about 200 micromolar in this assay. When testing inhibitory compounds, a substrate concentration of about 1.0 micromolar is preferred. Test compounds diluted in DMSO are added to the reaction mixture, with a final DMSO concentration of 5%. Controls also contain a final DMSO concentration of 5%. The concentration of beta secretase enzyme in the reaction is varied, to give product concentrations with the linear range of the ELISA assay, about 125 to 2000 picomolar, after dilution.

The reaction mixture also includes 20 millimolar sodium acetate, pH 4.5, 0.06% Triton X100, and is incubated at 37 degrees C for about 1 to 3 hours. Samples are then diluted in assay buffer (for example, 145.4 nanomolar sodium chloride,

9.51 millimolar sodium phosphate, 7.7 millimolar sodium azide, 0.05% Triton X405, 6g/liter bovine serum albumin, pH 7.4) to quench the reaction, then diluted further for immunoassay of the cleavage products.

5        Cleavage products can be assayed by ELISA. Diluted samples and standards are incubated in assay plates coated with capture antibody, for example, SW192, for about 24 hours at 4 degrees C. After washing in TTBS buffer (150 millimolar sodium chloride, 25 millimolar Tris, 0.05% Tween 20, pH 7.5),  
10    the samples are incubated with streptavidin-AP according to the manufacturer's instructions. After a one hour incubation at room temperature, the samples are washed in TTBS and incubated with fluorescent substrate solution A (31.2 g/liter 2-amino-2-methyl-1-propanol, 30 mg/liter, pH 9.5). Reaction  
15    with streptavidin-alkaline phosphate permits detection by fluorescence. Compounds that are effective inhibitors of beta-secretase activity demonstrate reduced cleavage of the substrate as compared to a control.

#### 20        **Example D**

##### Assays using Synthetic Oligopeptide-Substrates

Synthetic oligopeptides are prepared that incorporate the known cleavage site of beta-secretase, and optionally detectable tags, such as fluorescent or chromogenic moieties.  
25    Examples of such peptides, as well as their production and detection methods are described in U.S. Patent No: 5,942,400, herein incorporated by reference. Cleavage products can be detected using high performance liquid chromatography, or fluorescent or chromogenic detection methods appropriate to  
30    the peptide to be detected, according to methods well known in the art.

By way of example, one such peptide has the sequence SEVNL-DAEF [SEQ ID NO: 8], and the cleavage site is between residues 5 and 6. Another preferred substrate has the

sequence ADRGLTTRPGSGLTNIKTEEISEVNL-DAEF [SEQ ID NO: 9], and the cleavage site is between residues 26 and 27.

These synthetic APP substrates are incubated in the presence of beta-secretase under conditions sufficient to result in beta-secretase mediated cleavage of the substrate. Comparison of the cleavage results in the presence of the compound inhibitor to control results provides a measure of the compound's inhibitory activity.

#### Example E

##### Inhibition of Beta-Secretase Activity - Cellular Assay

An exemplary assay for the analysis of inhibition of beta-secretase activity utilizes the human embryonic kidney cell line HEKp293 (ATCC Accession No. CRL-1573) transfected with APP751 containing the naturally occurring double mutation Lys651Met52 to Asn651Leu652 (numbered for APP751), commonly called the Swedish mutation and shown to overproduce A beta (Citron et al., 1992, Nature 360:672-674), as described in U.S. Patent No. 5,604,102.

The cells are incubated in the presence/absence of the inhibitory compound (diluted in DMSO) at the desired concentration, generally up to 10 micrograms/ml. At the end of the treatment period, conditioned media is analyzed for beta-secretase activity, for example, by analysis of cleavage fragments. A beta can be analyzed by immunoassay, using specific detection antibodies. The enzymatic activity is measured in the presence and absence of the compound inhibitors to demonstrate specific inhibition of beta-secretase mediated cleavage of APP substrate.

#### Example F

##### Inhibition of Beta-Secretase in Animal Models of AD

Various animal models can be used to screen for inhibition of beta-secretase activity. Examples of animal models useful in the invention include, but are not limited

to, mouse, guinea pig, dog, and the like. The animals used can be wild type, transgenic, or knockout models. In addition, mammalian models can express mutations in APP, such as APP695-SW and the like described herein. Examples of transgenic non-human mammalian models are described in U.S. Patent Nos. 5,604,102, 5,912,410 and 5,811,633.

PDAPP mice, prepared as described in Games et al., 1995, *Nature* 373:523-527 are useful to analyze *in vivo* suppression of A beta release in the presence of putative inhibitory compounds. As described in U.S. Patent No. 6,191,166, 4 month old PDAPP mice are administered compound formulated in vehicle, such as corn oil. The mice are dosed with compound (1-30 mg/ml; preferably 1-10 mg/ml). After time, e.g., 3-10 hours, the animals are sacrificed, and brains removed for analysis.

Transgenic animals are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Control animals are untreated, treated with vehicle, or treated with an inactive compound. Administration can be acute, i.e., single dose or multiple doses in one day, or can be chronic, i.e., dosing is repeated daily for a period of days. Beginning at time 0, brain tissue or cerebral fluid is obtained from selected animals and analyzed for the presence of APP cleavage peptides, including A beta, for example, by immunoassay using specific antibodies for A beta detection. At the end of the test period, animals are sacrificed and brain tissue or cerebral fluid is analyzed for the presence of A beta and/or beta-amyloid plaques. The tissue is also analyzed for necrosis.

Animals administered the compound inhibitors of the invention are expected to demonstrate reduced A beta in brain tissues or cerebral fluids and reduced beta amyloid plaques in brain tissue, as compared with non-treated controls.



**Example G****Inhibition of A Beta Production in Human Patients**

Patients suffering from Alzheimer's Disease (AD) demonstrate an increased amount of A beta in the brain. AD patients are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Administration is repeated daily for the duration of the test period. Beginning on day 0, cognitive and memory tests are performed, for example, once per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as analyzed by changes in one or more of the following disease parameters: A beta present in CSF or plasma; brain or hippocampal volume; A beta deposits in the brain; amyloid plaque in the brain; and scores for cognitive and memory function, as compared with control, non-treated patients.

**Example H****Prevention of A Beta Production in Patients at Risk for AD**

Patients predisposed or at risk for developing AD are identified either by recognition of a familial inheritance pattern, for example, presence of the Swedish Mutation, and/or by monitoring diagnostic parameters. Patients identified as predisposed or at risk for developing AD are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Administration is repeated daily for the duration of the test period. Beginning on day 0, cognitive and memory tests are performed, for example, once per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as analyzed by changes in one or more of the following disease parameters: A beta present in CSF or

plasma; brain or hippocampal volume; amyloid plaque in the brain; and scores for cognitive and memory function, as compared with control, non-treated patients.

#### CHEMISTRY EXAMPLES

The following abbreviations may be used in the Examples:

EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide or the hydrochloride salt);

DIEA (diisopropylethylamine);

PyBOP (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate);

HATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate);

DCM (dichloromethane).

The following detailed examples and Schemes describe how to prepare various compounds of the invention and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques.

#### General Procedures

##### Synthesis of amides using HBTU

Coupling of the amine 8 (synthesis discussed in Example 1 below) with carboxylic acids is generally performed using the HBTU coupling method. A carboxylic acid is dissolved in anhydrous DMF (3mL/mmol) and 3 equivalents of PS-DIEA (polystyrene diisopropylethylamine, 3.5 meq/gram, Argonaut Corporation) is added followed by HBTU (1.1 equivalent relative to the acid). The reaction vessel is closed and the suspension is agitated. After 30 minutes one equivalent of the amine 8 in DMF solution is added and the reaction mixture is agitated overnight. Scavenger resins are added (5


equivalents each of Argonaut MP-carbonate and Argonaut PS-trisamine) are and the contents agitated for 5 hours. (Additional DMF is added if needed to improve mixing.) The reaction mixture is filtered and the solids are washed with 5 DMF. The solvent is evaporated from the combined filtrate and and the product is purified by chromatography.

#### Synthesis of amides using HATU

The coupling of carboxylic acids with secondary amines 10 such as 36 (discussed below) is performed using a procedure similar to that described in the previous paragraph, except that HATU is used in place of HBTU.

#### General Procedure for removal of BOC protecting groups 15 with 4 N HCl in dioxane

The Boc-protected compound is dissolved in at least 5 molar equivalents of 4N HCl in dioxane. The mixture is allowed to stand for two hours, and the solvent is then evaporated. The residue is dissolved in methanol and the 20 methanol is evaporated. The residue is redissolved in methanol and the methanol is evaporated again. The vial containing the product is then placed in a high vacuum dessicator (with NaOH pellets) overnight to yield the deprotected amine in the form of its hydrochloride salt. The 25 amine hydrochloride salt is optionally converted to the free base prior to use in amide coupling reactions. This is accomplished by partitioning it between ethyl acetate and a large excess of saturated aqueous potassium carbonate solution, washing the organic extracts with brine, and then 30 drying the solution over magnesium sulfate. The free base is then isolated by evaporation of the solvent.

**EXAMPLE 1: Synthesis of compound 8 in Scheme 4.****8**

**Intermediate 2:** To a 20 °C stirred suspension of hydroxyl  
5 compound 1 (53 mmol) in dichloromethane (300 mL) is added  
methanesulfonyl chloride (103 mmol) followed by the addition  
of diisopropylethylamine (200 mmol). The reaction mixture is  
stirred at room temperature for 1 h. The solution is then  
washed with 1 N potassium bisulfate followed by water and  
10 saturated sodium chloride. The organic phase is dried (sodium  
sulfate) and concentrated to provide mesylate 2.

**Intermediate 3:** To a stirred suspension of mesylate 2 (53  
mmol) in toluene (400 mL) is added diisopropylethylamine (104  
15 mmol). The reaction mixture is heated at 120 °C for 15 h. The  
reaction mixture is cooled to room temperature and 1 N KH<sub>2</sub>PO<sub>4</sub>  
(400 mL) is added. The organic layer is separated and the  
aqueous phase is extracted with ethyl acetate. The combined  
organic layers are washed with 1 N KH<sub>2</sub>PO<sub>4</sub>, water, and saturated  
20 sodium chloride, dried (sodium sulfate), filtered, and  
concentrated under reduced pressure. Purification by flash  
column chromatography provides oxazolidinone 3.

**Intermediate 4:** To a stirred solution of oxazolidinone 3  
25 (42 mmol) in tetrahydrofuran (100 mL) is added a suspension of  
sodium hydride (50 mmol) in tetrahydrofuran (80 mL). The  
resulting suspension is then treated with di-*t*-  
butyldicarbonate (48.9 mmol). The reaction mixture is stirred  
at room temperature for 1 h, methanol (90 mL) is added  
30 followed by the addition of a lithium hydroxide (214 mmol)

solution in water (90 mL). The reaction mixture is stirred at room temperature for 1 h. Aqueous 1 N  $\text{KH}_2\text{PO}_4$  (100 mL) and ethyl acetate (200 mL) were added. The organic phase is washed with 1 N  $\text{KH}_2\text{PO}_4$ , water, and saturated sodium chloride, dried (sodium sulfate), filtered, and concentrated under reduced pressure. Purification by flash column chromatography provides (4).

**Intermediate 5:** To a stirred solution of epoxide 4 (6.7 mmol) in N,N-dimethylformamide (20 mL) is added potassium cyanide (10.0 mmol) and water (2 mL). The reaction mixture is stirred at room temperature for 20 h and then quenched with 1 N  $\text{KH}_2\text{PO}_4$  (20 mL). The mixture is extracted with ethyl acetate (3 x 50 mL). The combined extracts are washed with saturated sodium chloride, dried (sodium sulfate), filtered, and concentrated under reduced pressure. Purification by flash column chromatography provides cyanohydroxy compound 5.

**Intermediate 6:** To a stirred solution of the Boc-protected amine 5 (1.23 mmol) in dioxane (2 mL) is added hydrogen chloride (solution in dioxane, 12 mmol). The reaction mixture is stirred at room temperature for 4 h and then diluted with ethyl ether (50 mL). The resulting precipitate is collected by filtration to provide amine hydrochloride 6.


25

**Intermediate 7:** To a stirred solution of acid 12 (1.23 mmol), diisopropylethylamine (1.3 mmol), and HBTU (1.35 mmol) in methylene chloride (8 mL) is added a mixture of 6 (1.3 mmol) and diisopropylethylamine (1.3 mmol) in methylene chloride (8 mL). The reaction mixture is stirred at room temperature for 16 h and concentrated under reduced pressure. The residue is diluted with ethyl acetate (50 mL), washed with saturated sodium bicarbonate, saturated sodium chloride, dried (sodium sulfate), filtered, and concentrated under reduced

pressure. Purification by flash column chromatography provides 7.


**Intermediate 8:** A mixture of cyano compound 7 (0.72 mmol) and platinum oxide (50 mg) in acetic acid (10 mL) is subjected to an atmosphere of hydrogen at 50 psi for 20 h. The mixture is filtered through diatomaceous earth and washed with methanol, and the filtrate is concentrated under reduced pressure. Purification by flash column chromatography provides 8.

**EXAMPLE 2: Synthesis of compound 9 in Scheme 4**




A mixture of amine 8 (0.028 mmol) and triethylamine (14 uL) in 0.5 mL of tetrahydrofuran is treated with 3.4 uL of propanesulfonyl chloride. The mixture is agitated for 6.5 hours and then cooled to 0 °C. An additional 14 uL of triethylamine and 1 uL of propanesulfonyl chloride is added. The mixture is allowed to warm to 25 °C and stirred an additional 16 hours. The solvent is evaporated and the product is isolated by reverse phase HPLC to provide 9.

**EXAMPLE 3: Synthesis of compound 10 in Scheme 4**




A solution of 0.028 mmol of amine 8 in 0.5 mL of tetrahydrofuran is treated with a solution of 2.6 uL of propylisocyanate in 100 uL of tetrahydrofuran. After agitating for 24 hours, a solution of 0.7 uL of propylisocyanate in 25 uL of tetrahydrofuran is added. The mixture is agitated for another 2 hours and then the solvent is evaporated in a stream of nitrogen. Purification by reverse phase HPLC gives 10.

**EXAMPLE 4: Synthesis of compound 11 in Scheme 4**



A solution of 0.028 mmol of primary amine 8 in 0.5 mL of tetrahydrofuran is treated with 5.7 uL of butyric anhydride. After 23 hours, 60 mg of trisamine resin (Argonaut Technologies) is added and the mixture is agitated an additional 6 hours. The mixture is filtered and the solvent is evaporated to give compound 11.

**Example 5: Synthesis of compound 38 in Scheme 5**




**38.**

Amine 8 is acylated with 3-(1-(tert-butoxycarbonyl)-piperidin-4-yl)propanoic acid under HBTU conditions to give amide 35. The BOC group is cleaved with HCl in dioxane to give amine hydrochloride salt 36. This hydrochloride salt is acylated with N-Boc-6-aminohexanoic acid to give compound 37.

The Boc group of 37 is in turn cleaved with HCl in dioxane to give compound 38.


**Example 6: Synthesis of compound 40 in Scheme 6**



**40.**

To 0.084 mmol of 8 in 0.25 mL of methanol is added 12 mg of methyl N-cyanobutyrimidate (*J. Med. Chem.* vol 36, 2253-2265). After 15 hours the solvent is evaporated and the residue is purified by reverse-phase HPLC to give 40.


**Example 7: Synthesis of compound 41 in Scheme 6**



**41.**

To 0.05 mmol of 8 in 0.25 mL of isopropanol is added 10 mg of 2-chloro-4-(trifluoromethyl)pyrimidine and 19  $\mu$ L of diisopropylethylamine. The mixture is stirred at 50 °C for 24 hours and then it is filtered. The product is isolated by reverse-phase HPLC.




**Example 8: Synthesis of compound 48 in Scheme 7**COMPOUND 42

5 To NaOEt (10 mmol, 3.3mL of 3N NaOEt in EtOH) in EtOH (27 mL) was added diethylmalonate (12 mmol, 1.8 mL). The mixture was stirred at rt for 15 min, then a solution of 4 (3 g, 10 mmol) in EtOH (17 mL) was added. The mixture was stirred at rt overnight. The reaction mixture was concentrated to a residue and 10% citric acid was added. The mixture was extracted with EtOAc twice. The organic layers were combined and washed with sat. NaHCO<sub>3</sub> and Brine. It was dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated to a residue and purified by column chromatography eluting with 10 - 33% ethyl acetate in hexanes to afford 42 (2.8g, 6.8 mmol, 68% yield) as a white solid. LC/MS: 436.2 (M + Na).

10

15

COMPOUND 43


20

To NaOEt (10 mmol, 2.56 mL of 3N NaOEt in EtOH) in EtOH (10 mL) was added 42 (2.8g, 6.7 mmol) in EtOH (36 mL) at 50 °C. Then CH<sub>3</sub>I (0.78 mL) was added to the mixture. The mixture was stirred at at 50 °C for 4 hr. The reaction mixture was quenched with 1N NaH<sub>2</sub>PO<sub>4</sub> and concentrated to remove EtOH. Then it was partitioned between EtOAc and H<sub>2</sub>O. The organic layers


25

were combined and washed with H<sub>2</sub>O and brine, then dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated to a solid. This solid was dissolved in THF (70 mL) and LiOH (1M in H<sub>2</sub>O, 35 mL) was added. The resulting mixture was stirred at room  
5 temperature for 1 day. It was quenched with 10% citric acid and concentrated to remove THF. It was extracted with EtOAc. The organic phase was dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated to a residue and toluene (50 mL) was added and the mixture was refluxed for 2 days. It was  
10 concentrated and the residue was purified by normal phase HPLC to afford two isomers. The desired isomer 43 (0.6 g, 1.7 mmol, 25%) was collected as a white solid. LC/MS: 378.1 (M+Na)


15

COMPOUND 44

To 43 (0.4g, 1.13 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added TFA (1  
20 mL) at rt. The mixture was stirred at rt for 1 hr. Then it was concentrated to a residue and was dissolved in EtOAc. It was washed with sat. NaHCO<sub>3</sub> and H<sub>2</sub>O. The organic layer was dried and filtered. The filtrate was concentrated to afford 44 (280 mg, 100%) as an oil: LC/MS: 256.1.

COMPOUND 45


To 44 (1.1 mmol) in DMF (6 mL) is added acid 12 (1.5 mmol), followed by HOBT (1.46 mmol) and TEA (0.4 mL). Then  
 5 EDC (1.46 mmol) is added to the stirring mixture at room temperature. The mixture is stirred at room temperature overnight. It is quenched with 1N NaH<sub>2</sub>PO<sub>4</sub> and extracted with EtOAc. The organic layer is washed with 1N NaH<sub>2</sub>PO<sub>4</sub>, sat. NaHCO<sub>3</sub>, H<sub>2</sub>O and Brine. It is dried with MgSO<sub>4</sub> and filtered.  
 10 The filtrate is concentrated to a solid which contains 45.

COMPOUND 46

15 To crude 45 (0.9 mmol) in DME (6 mL) is added LiOH (1N, 6 mL). The mixture is stirred at rt for 3 hr. Then 10% citric acid is added and it is extracted with EtOAc. The organic layer is washed with brine, dried with MgSO<sub>4</sub> and filtered. The filtrate is concentrated to a residue and used immediately.  
 20 Thus the residue is dissolved in DMF (6 mL), then imidazole (0.75 g) and TBDMSCl (1.93 mL solution of 2.86 M in toluene) is added. The mixture is stirred at rt for a day. MeOH (20 mL) is added and the mixture is stirred at rt overnight. The mixture is concentrated to a residue and partitioned between


10% citric acid and EtOAc. The organic layer is washed with brine. It is dried with  $\text{MgSO}_4$  and filtered. The filtrate is concentrated to a residue and purified by column chromatography to afford **46**.

# 5 COMPOUND 47



To **46** (0.19 mmol) in  $t$ -butanol (5 mL) is added TEA (0.19 mmol) and DMAP (2 mg). Then DPPA (0.19 mmol) is added to the mixture and the mixture is heated to  $100^\circ\text{C}$  for 5 hr. The reaction mixture is partitioned between EtOAc and  $\text{H}_2\text{O}$ . The organic layer is dried with  $\text{MgSO}_4$  and filtered. The filtrate is concentrated to a residue which is purified by prep TLC to afford **47**.


# 15 COMPOUND 48







To **47** (0.056 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 mL) is added HCl (4 M in dioxane 1 mL). The mixture is stirred at rt for 1 hr. Then the mixture is concentrated and the residue is dissolved in DMF (300  $\mu\text{L}$ ). To this solution is added valeric acid (0.084 mmol), HOBT (0.084 mmol) and TEA (0.17 mmol). Then EDC (0.084



mmol) is added to the mixture and the mixture is stirred at rt for 2 hr. The resulting mixture is extracted between 10% citric acid and EtOAc. The organic layer is washed with sat. NaHCO<sub>3</sub> and brine. It is dried with MgSO<sub>4</sub> and filtered. The filtrate is concentrated to a residue and purified by HPLC to afford 48.

The following compounds are prepared essentially according to the procedures described in the schemes, charts, examples and preparations set forth herein.


Ex. No.	Name/Structure
9	 <p><i>N</i>-[(3<i>S</i>,4<i>S</i>)-4-(acetamido)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-4-methylpentanamide</p>
10	<i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetamido)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-4-methylpentanamide
11	<i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetamido)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-4-methylpentanamide
12	2-(acetamido)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(4-methylpentanoyl)amino]- <i>L</i> -threo-hexitol


13	 <p>2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(4-methylpentanoyl)amino]-L-threo-heptitol</p>
14	<p><i>N</i>-[(3<i>S</i>,4<i>S</i>)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-4-phenylbutanamide</p>
15	 <p><i>N</i>-[(3<i>S</i>,4<i>S</i>)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-4-phenylbutanamide</p>
16	 <p><i>N</i>-[(3<i>S</i>,4<i>S</i>)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-4-phenylbutanamide</p>
17	<p>2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(4-phenylbutanoyl)amino]-L-threo-hexitol</p>

18	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(4-phenylbutanoyl)amino]-L-threo-heptitol
19	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-2-(benzyloxy)acetamide
20	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-2-(benzyloxy)acetamide
21	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-2-(benzyloxy)acetamide
22	2-(acetylamino)-5-{[(benzyloxy)acetyl]amino}-1,2,4,5-tetradexo-1-(3,5-difluorophenyl)-L-threo-hexitol
23	2-(acetylamino)-5-{[(benzyloxy)acetyl]amino}-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-L-threo-heptitol
24	 <p>N-[(3<i>S</i>,4<i>S</i>)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-3-cyclopentylpropanamide</p>
25	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-3-cyclopentylpropanamide
26	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-3-cyclopentylpropanamide



27	 <p>2-(acetylamino)-5-[(3-cyclopentylpropanoyl)amino]-1,2,4,5-tetrahydroxy-1-(3,5-difluorophenyl)-L-threo-hexitol</p>
28	<p>2-(acetylamino)-5-[(3-cyclopentylpropanoyl)amino]-1,2,4,5,6-pentahydroxy-1-(3,5-difluorophenyl)-L-threo-heptitol</p>
29	 <p>N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-2-ethoxyacetamide</p>
30	<p>N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-2-ethoxyacetamide</p>
31	<p>N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-2-ethoxyacetamide</p>
32	<p>2-(acetylamino)-1,2,4,5-tetrahydroxy-1-(3,5-difluorophenyl)-5-[(ethoxyacetyl)amino]-L-threo-hexitol</p>
33	<p>2-(acetylamino)-1,2,4,5,6-pentahydroxy-1-(3,5-difluorophenyl)-5-[(ethoxyacetyl)amino]-L-threo-heptitol</p>



34	<i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]-2-propoxyacetamide
35	<i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]-2-propoxyacetamide
36	<i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]-2-propoxyacetamide
37	2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(propoxyacetyl)amino]- <i>L</i> -threo-hexitol
38	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(propoxyacetyl)amino]- <i>L</i> -threo-heptitol
39	 <p>(3<i>E</i>)-<i>N</i>-[(3<i>S</i>,4<i>S</i>)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]hex-3-enamide</p>
40	(3 <i>E</i> )- <i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]hex-3-enamide
41	(3 <i>E</i> )- <i>N</i> -[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]hex-3-enamide
42	2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(3 <i>E</i> )-hex-3-enoylamino]- <i>L</i> -threo-hexitol

43	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(3E)-hex-3-enoylamino]-L-threo-heptitol
44	(3E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]pent-3-enamide
45	(3E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]pent-3-enamide
46	(3E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]pent-3-enamide
47	2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(3E)-pent-3-enoylamino]-L-threo-hexitol
48	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(3E)-pent-3-enoylamino]-L-threo-heptitol
49	(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]hex-2-enamide
50	(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]hex-2-enamide
51	 <p>(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]hex-2-enamide</p>

52	2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(2E)-hex-2-enoylamino]-L-threo-hexitol
53	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(2E)-hex-2-enoylamino]-L-threo-heptitol
54	(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]pent-2-enamide
55	(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]pent-2-enamide
56	(2E)-N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]pent-2-enamide
57	2-(acetylamino)-1,2,4,5-tetradecoxy-1-(3,5-difluorophenyl)-5-[(2E)-pent-2-enoylamino]-L-threo-hexitol
58	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-[(2E)-pent-2-enoylamino]-L-threo-heptitol
59	N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]pentanamide
60	N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]pentanamide
61	N-[(3S,4S)-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]pentanamide

62	 <p>2-(acetylamino)-1,2,4,5-tetradeoxy-1-(3,5-difluorophenyl)-5-(pentanoylamino)-L-threo-hexitol</p>
63	 <p>2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-(pentanoylamino)-L-threo-heptitol</p>
64	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-methylpentyl]hexanamide
65	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-1-ethyl-3-hydroxypentyl]hexanamide
66	N-[(3 <i>S</i> ,4 <i>S</i> )-4-(acetylamino)-5-(3,5-difluorophenyl)-3-hydroxy-1-propylpentyl]hexanamide
67	2-(acetylamino)-1,2,4,5-tetradeoxy-1-(3,5-difluorophenyl)-5-(hexanoylamino)-L-threo-hexitol
68	2-(acetylamino)-1,2,4,5,6-pentadeoxy-1-(3,5-difluorophenyl)-5-(hexanoylamino)-L-threo-heptitol

The invention has been described with reference to various specific and preferred embodiments and techniques.

However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.